THE MEDICAL NEWS.

A WEEKLY JOURNAL OF MEDICAL SCIENCE.

Vol. LXII.

SATURDAY, JUNE 24, 1893.

No. 25.

ORIGINAL ARTICLES.

ICE IN THETREATMENT OF ACUTE PNEUMONIA. A Collective Report.

BY THOMAS J. MAYS, M.D.,

PROPESSOR OF DISEASES OF THE CHEST IN THE PHILADELPHIA POLY-CLINIC, VISITING PHYSICIAN TO THE RUSH HOSPITAL FOR CONSUMPTIVES. BTC.

WHATEVER its nature may be, it is quite certain that no other disease has elicited a greater number of conflicting opinions concerning its treatment than has croupous pneumonia. Forty years ago bleeding and blistering were regarded as its specifics; but these are now, and for the last twenty years have been, scarcely thought of in this connection. In the meantime hot poultices, aconite, veratrum viride, digitalis, quinine, etc., took their places; vet it is not too much to say that these have all led to disappointment and have come to grief in the retort of clinical experience, and that finally the profession has gravitated to the conviction that the disease is self-limited in duration, and that hence all efforts to control its course are fruitless, if not actually harmful.

To be thus compelled to stand before a disease and acknowledge our helplessness and impotency is, to say the least, an unenviable position, but I must confess that until I became familiar with the value of local cold applications in pneumonia I was in entire accord with this idea. Since then I may say that I am able to approach a case of pneumonia with a greater degree of assurance—not with the feeling, however, that we possess a specific, but with the confidence that we have in cold an agency with which we are able to impress and circumvent the severity of the pneumonic process. I believe that cold properly applied will affect the death-rate of pneumonia as profoundly as it has affected that of typhoid fever, and, although I do not expect a rapid introduction of this measure, on account of a deeprooted prejudice which exists against the use of cold in almost all internal diseases, I trust that the evidence which is herewith submitted will serve to commend it to the serious attention of the profes-

Under the titles "Can Croupous Pneumonia be Aborted?" and "Ice in the Treatment of Croupous Pneumonia," I contributed two papers to THE MEDICAL News of September 24, 1892, and January 21, 1893, respectively, in which are related

three cases of pneumonia that were treated principally with applications of ice to the chest. After the appearance of the first paper I instituted a collective investigation on a small scale by sending a number of circulars¹ to various members of the profession, inviting a trial of the ice-treatment, and as a result of this inquiry I submit the following abstracts of cases which have been kindly returned to me, and also of those of other cases which I have collected from the literature on the subject.

Case I.2—M., aged thirty-six years, was attacked with right basic pneumonia March 10, 1892, when the temperature reached 105°, pulse 112, and respiration 26. Ice was applied over the affected area until the 16th inst., when crisis followed the hypodermatic injection of ¼ of a grain of pilocarpine, and the temperature fell to normal. Recovery was retarded by fibrinous pleurisy. The patient was able to sit up on April 11th.

Case II.— F., aged seventy-four years, had pneumonia of the right base, associated with delirium. Ice was applied over the affected area from the 6th to the 10th of March, 1892, when the crisis occurred and the ice-treatment was discontinued. The temperature and pulse records are lost, but the im-

provement was prompt and rapid.

CASE III.—M., four years of age, was first seen March 6, 1892, when the child was intensely delirious and quite insensible. Pneumonia of the right base existed. Ice was applied from March 7th to the 13th, when lysis took place. There was no pulse or temperature record kept, but it was "a very severe case, in which the cold acted like a charm."

Case IV.—M., fourteen years old, had pneumonia of both bases, the consolidation extending above the nipple on the right side. Ice was applied around the whole chest from the 22d to the 29th of March, 1892. At the beginning of the application the temperature was 105°, pulse 100, and respiration 28. Crisis occurred on the 28th, when the temperature became normal.

CASE V.—M. aged two years, had pneumonia, most severely at the left base, and was very delirious. Ice was applied March 12th, when the temperature was 101°, pulse 150, respiration 40. The ice-applications were continued until the 14th, when crisis took place, and temperature became normal. "I think this can be taken as an aborted case."

Case VI.—F., aged twenty-three, had a chill January 7, 1892, which was followed by pneumonia

Similar circulars will be cheerfully sent by the author to any one who may apply for them.

² The first eleven cases are reported by Dr. W. Fred. Jackson, of Brockville, Ontario,

implicating the whole of the right lung. On January 8th, when the temperature was 105.2°, the ice was applied, and by next morning the temperature had fallen to 99°, and the next day it was normal. In order to satisfy the patient the ice was continued until the 11th. He also has an organic heart-lesion, which aggravated his condition very much.

CASE VII.—M., aged thirty-four, had pneumonia of the right base. When first seen, January 12th, he was insensible, the temperature was 103°, and he had rusty and bloody expectoration. The ice was applied on this day, and the next morning the temperature had fallen to 100°, but owing to some objection it was taken off, and by evening the temperature had risen to 103.2°. The ice was promptly reapplied, and the following day the temperature reached 100°; a day later crisis occurred and the temperature fell to normal.

Case VIII.—F., aged sixty-five, had pneumonia of the whole of the right lung. Ice was applied to this side March 16th, when the temperature was 103°; next morning it had fallen to 101°. Owing to prejudice, the applications were discontinued until the morning of the 22d, when she was no better, and the temperature had again risen to 103°. The ice was then reapplied. On the evening of this day the temperature was 101°; the morning and evening of the 23d it was 100° and 99.6°, and crisis occurred on the 24th, when the temperature became normal.

Case IX.—M, aged forty, had pneumonia of the upper half of the left side and right base. The sputum was bloody. On the morning of February 5, 1892, the date of his initial chill, the ice was applied; his temperature was 104.4°, and in the evening 102°. On the 6th, the morning record was 100°, the evening 99°. On the 7th, crisis supervened and the temperature became normal. The pain which this person experienced was also quickly relieved by the ice. The disease was "aborted in the congestive stage."

CASE X — F., aged thirty-five, had pleuro-pneumonia of the left base. The initial chill was on February 5th, and ice was applied over the left lung on the morning of the 6th, when the temperature was 101°. On the evening of the same day, it was 100°; on the 7th, morning 99.4°, evening 99°. On the 8th, crisis took place, and the temperature was 98°.

Case XI.—M., aged thirteen, had pneumonia of both bases. Ice was applied around the chest on March 10, 1892, when the temperature was 105°; on the 11th, the record was 103°, and on the 12th, 102°. On the 13th crisis occurred, with a thermometric record of 98°. "A very severe case in which the effect of the cold was very marked." (All of Dr. Jackson's cases received liq. ammon. acetat. 3ij. and spt. ether. nit. 3j. every hour, well diluted.)

Case XII. (Reported by Dr. E. W. Blackburn, Stahlstown, Pa.)—The patient was a male, a periodical drinker, aged forty-one, and had pneumonia over the whole of the right lung. The initial chill was on November 10, 1892, following a three days' alcoholic debauch, and the man was first seen on the following day, when his temperature was 103.2°. The

ice was then applied. On the 12th, the temperature was 101°; on the 13th, 99°; on the 14th, lysis occurred, with a normal temperature. This patient also received $\frac{1}{10}$ grain of strychnine every four hours.

CASE XIII.—(The following four cases are reported by Dr. J. C. Knipe, Jewish Hospital, Philadelphia.) The patient, a male, aged thirty-seven, an alcoholic, was taken with a chill, which was followed by pneumonia of the left base, on March II, 1893, two days before admission to the hospital. He was suffering from diarrhea, lead-poisoning, and while in the hospital had an attack of delirium tremens. His temperature ranged from 102° to 103°; the ice was applied for three days, and death took place on the fifth day in the hospital.

Case XIV.—A male, aged twenty-eight, with pneumonia of the left lung, was admitted October 17th, although he had a chill two days previously. He was delirious, and had rusty and bloody expectoration. The ice was applied over the affected area for five days. His temperature on admission was 104°, afterward from 100° to 102°. He died October 22d. Before this attack he had been suffering from chronic pneumonia of both bases.

CASE XV.—A male, aged seventeen, had pneumonia of the right base, December 27, 1892. On admission, when the ice was applied, his temperature was 103°, and for the next twelve days, during which the ice was continued, it ranged from 101° to 98°, and disease terminated by lysis at the end of that time.

CASE XVI.—A male, aged twenty one, was seized with pneumonia of the right base February 2, 1893. The ice was applied for six days over the affected area. The temperature remained at 104° and 105.6° until midnight of the second day of the disease, when it dropped to 99.4°. Shortly after this time the patient had another chill, and on the third day, in the morning, the temperature rose to 105°, and in the evening, after a profuse sweat, it fell to 98°; from this time it fluctuated between 104° and 98°, until the seventh day, when it became normal, and remained so. This patient also received aconite, morphine, and ammonium carbonate.

CASE XVII.—(The next two cases are reported by Dr. John A. Fell, of Doylestown, Pa.)-M., aged sixty, had the initial chill, December 20, 1892, and from this time to the afternoon of the 28th, the temperature varied from 102° to 103.5°. On the last date the ice was applied over the lower half of the right lung, the seat of the pneumonia, and at noon of January 1st the temperature was normal, when it was removed. He also received 1 grain of strychnine every four hours until January 1st. This patient also had an old-standing aortic regurgitation. At first he objected to the use of the ice, but it gave him such marked relief from his distressed breathing that he became much pleased with it. He also received two ounces of fresh beef-juice and a glass of milk alternately every two hours.

Case XVIII.—F., aged seventeen, had pneumonia of the left lung. For the first three days the tinctures of aconite, belladonna, and veratrum viride were used in large doses, and mustard was applied to the affected side, without any apparent effect. Her

temperature was about 103°, and pulse 120. On the fourth day the ice-bags were applied, and in less than twenty-four hours the temperature had fallen to 99°, and the patient experienced a sense of comfort which she had not felt before. Strychnine, quinine, and morphine, were also used. "Her recovery was rapid and complete after using the ice."

CASE XIX. (Reported by Dr. A. Cooper, Brookline, Mass.)-A male, aged thirty-one, had the initial chill on the evening of January 31, 1893, and the expectoration was of a bloody and mucous character from this date. Both bases were involved. On the evening of February 1st the temperature was 102.5°; the next morning it was 104.25°, when the ice was applied. On the evening of this day it was 103°. On the third day it fell to 101°, and was almost normal on the fifth day. The ice was applied for seventy-two hours to the right, and for eighty-six hours to the left side, and the disease subsided by lysis. Dr. Cooper adds: "In six hours after the ice had been applied the temperature dropped from 104.25° to 101°. The ice had been on a very short time when pain and dyspnea ceased. My patient was better on the fifth day than was any case I have had under poultice or hot local treatment on the ninth or twelfth day.'

CASE XX. (Reported by Dr. G. H. Franklin, Hightstown, N. J.)—F., five years of age, had an initial chill February 17, 1893, followed by pneumonia of the entire right lung. From this date to the 20th the temperature rose from 102.5° to 104°, and she received the usual treatment for this disease. On the latter day ice was applied to the front and side of right lung, and was continued for seven days, during which time the temperature gradually returned to nearly a normal point (99°). Previously to the ice-treatment large doses of quinine had exerted no influence on the fever. The patient also received $\frac{1}{120}$ grain of strychnine every four hours. This was a case of broncho-pneumonia following measles.

Case XXI. (Reported by Dr. Eugene P. Bernardy, Philadelphia.)—M., aged twenty-five, was first seen September 15th, although the chill had occurred three days previously. The lower two-thirds of the right lung were involved. Expectoration was copious, rusty, and bloody. The patient was violently delirious; his temperature ranged from 102° to 104°. Ice was applied on the 16th, and was continued to the 19th On the evening of the last date active delirium set in, and the ice was discontinued. After the delirium was under control ice was again applied for three days longer, when the disease terminated by crisis. The patient also received 10 gr. of strychnine every two hours.

Case XXII. (Reported by Dr. Charles E. Woodward, West Chester, Pa.)—F., aged thirteen, with double pneumonia, had the initial chill March 20, 1893, accompanied by a temperature of 104°. The expectoration was bloody at first, afterward rusty. Crisis took place on the seventh day. Turpentine stupes were applied externally, and strychnine in 180 grain doses was given internally every four hours from the beginning. General sponging with cold

water (65°) was used throughout, and ice was applied only three times, and fifteen minutes each time. The superiority of the ice-bag over the cold sponging, was shown by the temperature-record. While the cold sponging reduced the temperature, which was about 104° for seven days, the reduction caused by it was not so pronounced as that which was caused even by the fleeting local application of ice. It is stated that the patient experienced great comfort from the use of the ice-bag.

CASE XXIII. (Reported by Dr. E. R. Snyder, Philadelphia.)-F., aged three years, had double pneumonia, the lower third of the left lung being involved at the beginning, and on the fifth day the lower part of the right lung became implicated. On the sixth day the temperature had risen to 105.4, and the child became unconscious. The whole chest was now enveloped in ice, and in six hours, when the temperature had fallen to 102.5°, the ice was removed. In the next six hours the temperature rose to 104°, when the ice was reapplied, and in one hour the thermometer registered 102.5°. After this the temperature gradually dropped to the normal point. There was good reason for believing that the ice saved the child's life. Strychnine was given hypodermatically in 1 grain doses, and croton oil was applied externally.

Case XXIV. (Reported by Dr. Walter Webb, Sharon Hill, Pa.)—A male, aged sixteen, had pneumonia of the entire left lung and lower portion of the right. The patient was seen first November 2, 1892, when the temperature was 105°, respiration about 60, and pulse 120. The parents objected to the ice when Dr. Webb first suggested it, but as no improvement in the boy's condition was observed, the parents permitted its use on November 6th, when the evening temperature was 105.2°, and the respiration 64. After this the fever gradually fell until the 14th, when it registered 102° in the evening. The ice was now withdrawn and the temperature rose a degree or more, and the boy complained of severe pain and greater difficulty in breathing. The ice was reapplied on the 17th, the temperature fell, and the patient expressed himself as being greatly relieved when the ice was reapplied. It was continued for two weeks, when the fever had disappeared.

The following six cases are my own:

CASE XXV.—F., twenty-five years of age, with double basic pneumonia, and with pleurisy on the left side, was first seen in the Polyclinic Hospital, November 24, 1891. She had a temperature of 103.4°, and ice-bags were applied over the affected area. The temperature began to decline immediately, and, excepting a slight rise on December 3d, sank to nearly a normal level on the eleventh day.

CASE XXVI.—F., thirty seven years old, was seen first in the Polyclinic Hospital, June 25, 1892, when she was suffering from pneumonia of the lower half of the right lung, and had a temperature of 103°. Ice was applied at once over the affected area, and by the following morning the temperature had descended to 98.8°, although in the afternoon it rose again to 101.5°. On the

morning of the third day the thermometer registered | 98.2°, and in the evening 100.8°. After this, with a few variations, the temperature became normal, and remained so.

Case XXVII.—M., fifteen years of age, had been having pneumonia of the lower two-thirds of his right lung for about six days before I saw him. On the evening of the second day his temperature was 104.8°, and cold-sponging was resorted to until the sixth day, when the temperature was reduced to 102.8° at 2.30 P.M. Ice-bags were now put over his right chest, and at 1 P.M. of the next day the temperature registered 98.2°, and at 8 P.M. 100°. On the following morning it was reduced to 97°, when the ice-bags were removed until reaction was assured. From this time on, the temperature, with a few exceptions, remained near the normal line, and the boy made a complete recovery.

CASE XXVIII.—A female, eighteen years of age, was first seen by me June 17, 1892, suffering with pneumonia of the whole of the right lung, and having a temperature of 102.8°. The whole right chest was surrounded with ice-bags on the third day of her disease, and by the next morning the tempera-ture had fallen three degrees (to 99.6°), and thereafter became gradually normal. Complete restora-

tion followed at once.

CASE XXIX.—A male, aged forty-five, an inveterate drinker for twenty years, was admitted into the Polyclinic Hospital, December 15, 1892, on the fifth day of a pneumonia involving the whole of the left lung, and with a temperature of 102.6°. The whole left chest was enveloped in ice, and on the morning of the eighth day the temperature was With the exception of a severe attack of delirium, which occurred on the ninth day, the man made an uninterrupted recovery.

CASE XXX.-M., aged thirty-four, had pneumonia of the whole left lung when first seen in the Polyclinic Hospital, March 26, 1893. His evening temperature was 103.8; ice was applied, and the the next morning the pyrexia had fallen to 100.4°. It remained near 101° until the 28th, when it rose to 102.4°. It was now found that the whole lower half of the opposite lung was involved. The man had several attacks of hemoptysis, his face was intensely flushed, and his breathing extremely difficult. Inhalation of oxygen relieved the distress somewhat. The patient was delirious, and had a tendency to coma. The ice was continued on both sides, and the temperature, except on several occasions, was kept between 101° and 102°. Resolu-tion soon followed in both lungs. Owing to the gravity of the condition convalescence was slow, but the man made a good recovery. The ice was continued for twenty days. In addition to the oxygen-inhalations, the patient received strychnine in large doses, by the mouth and hypodermatically, and was liberally fed and stimulated.

The next eighteen cases are reported by Dr. D. B. Lees in an article published in The Lancet, November 2, 1889:

CASE XXXI.-M., twenty-five years of age, had pleuro-pneumonia of the right base. On November 15, 1887, the fifth day of the disease, the temperature was 104.8°. The affected side was tightly strapped and an ice-bag applied over the strapping. temperature fell immediately, and was normal the

next morning, when the ice-bag was removed.

CASE XXXII.—M., aged twenty-six, with pneumonia of the left posterior base, was seen on the seventh day, when the temperature was about 104° and ice was applied. During the following night the temperature fell to 99°.

CASE XXXIII.-M., sixteen years old, with pneumonia of the right base, was seen on the fourth day, when his temperature was 104°. Snow was at once applied, and the temperarure fell immediately.

CASE XXXIV.—A child, three years old, had broncho-pneumonia of the base of the lung. The temperature was 104°, and the breathing was very rapid. The ice-bag was applied over the affected lung. Within twenty minutes of its application a distinct change for the better was observed. The breathing became much quicker, the temperature fell at once, and in a few days the patient was quite well.

CASE XXXV.-M., four years of age, had pneumonia of the left posterior base. The temperature was 104.5°, the pulse 140, the respiration 50. Ice-bags were applied over the implicated lung-area, and im-

mediate recovery followed.

CASE XXXVI.-F., four years old, suffered with pneumonia of the lower part of the right lung. When first seen the temperature was 104°, and within a few hours after the ice was applied it fell to 98°. However, it rose again to 101°, and continued at this point for about thirty-six hours, when it fell to

96.5°, and remained subnormal.

Case XXXVII.—F., thirty-eight years of age, had pneumonia over the left axillary base. woman was seen first on the fourth day of her illness, when her temperature was 103°. The ice-bag was applied, and by midnight her fever had fallen to 99.6°, and the ice-bag was removed. The temperature rose at once, and the next morning it was 101.5°, and in the evening 104°. No ice was again applied, but the temperature gradually fell, so that on the evening of the next day it reached 99.2°.

CASE XXXVIII.—M., nine years of age, had pleuro-pneumonia of the left base, and a temperature of 103°; ice was applied, and in thirty-six hours it had dropped to 99.4°, and the ice was removed. In eight hours it had risen again to 104°. Reapplication of the ice caused the temperature to sink to 98.4° in sixteen hours, after which it did not again

rise above 100°

CASE XXXIX.—M., twenty-six years old, had pneumonia of the left base. The man was seen on the third day, when his temperature was 103°. An ice-bag was applied, and the temperature fell promptly to 97.6°. On the morning of the fourth day the ice was removed, and the temperature rose When it reached 100° the ice-bag was replaced. The rise, however, continued up to 102°, from which point the temperature fell again to the normal on the fifth day.

CASE XL.-F., six and a half months old, had broncho-pneumonia of the root of the right lung, with a temperature of 105°. Twenty-four hours after the

application of the ice-bag the pyrexia fell to 101.4°. The ice was then removed, and the temperature rose to 103°, but fell again to 100.4°. A sudden rise to 105.6° was caused by an implication of the root of the opposite lung. The ice-bag being applied to the new focus of inflammation, the fever fell to 98.8° in nine hours, and after a few small transitory rises, re-

covery slowly occurred. CASE XLI.-F., fifteen years old, had an attack of broncho-pneumonia, with general bronchitis. There was dulness over the whole front of the left lung, with a temperature of 104.2°, on January 24, 1885, when the girl was first seen. Poultices were applied, and the next day she was worse, the tempera-ture having risen to 105°. Venesection—ten ounces—gave great and immediate relief. The temperature fell 1°, and on the evening of the next day (the third in the hospital) the child was worse than ever, her temperature reaching 105.6°. An ice-bag was now applied over the whole left front of the chest, and on the morning of the next day (the fourth) the thermometer registered 98°. In the afternoon, however, it rose again to 105.8°. Examination showed that the resonance over the left front was remarkably improved, but fresh crepitation and bronchial breathing had appeared in the axillary region. The ice-bag was shifted to this region, and again the temperature fell to 98°. It rose once more to 103°, and fell again to 98°. On the fourth day of the ice-treatment there was another rise to 104.2°, which proved to be due to the implication of the opposite apex. A second ice-bag was applied here, and in twenty-four hours the apex was normal, and the temperature had fallen to 96.7°. After this convalescence was uninterrupted.

CASE XLII.—M., thirteen years of age, had pneumonia of the left base. The boy was first seen on the fourth day of the disease, when the temperature was 105.8° Five grains of quinine were given, and the next morning the temperature was only 100.6°, but in the afternoon it rose to 104.2°, when an ice-bag was applied over the inflamed lung. The rise continued until 6 P.M., when 105.4° was reached, and thereafter there was a gradual abatement until the seventh day, when the temperature was 99°, and the ice-bag was removed.

CASE XLIII.—F., twenty years old, was first seen on the seventh day of an attack of pneumonia of the right base, at which time her temperature was 104.4°, and an ice-bag was applied. There was a fall of three degrees during the next few hours, but on the following morning the temperature stood again at 104°, and remained between 103° and 104° for three days. On the tenth day it suddenly fell to to 98°, and the ice was removed. On the eleventh and twelfth days the pyrexia rose to 102°. Examination now showed a fresh inflammation at the base of the opposite lung. The ice was applied to this spot, and in three days the physical signs had disappeared, and the woman was convalescent.

CASE XLIV.—M., twenty-five years old, was admitted on the fourth day of an attack of pneumonia of the right apex, and was seen first on the following day, with a temperature of 103.6°, and in a semi-maniacal condition. An ice-bag was ap-

plied to the chest, and on the evening of the eighth day the temperature dropped to 99°, and the ice was removed.

CASE XLV.-F., twenty years old, was seen first, with pneumonia of the right apex, on the fifth day, and with a temperature of 104°. By the next morning the ice had depressed the temperature to 100°, but during the day it rose to 102°. The respiration now rose to 74, and some cyanosis appeared. It was found that the lung in front and behind was dull to the base of the angle of the scapula. On the next day (the seventh) the temperature was only 102.5°, but the respiration had risen to 100, and the right lung was involved throughout. Both cheeks were markedly cyanotic, and the expectoration was of a "prune-juice" character. The whole right chest was enveloped in ice, and the patient received four ounces of brandy daily, and a mixture of ether and ammonia every four hours. On the eighth day the temperature was 101.5°, but the respiration ranged from 88 to 100. On the ninth and tenth days the temperature remained about the same, but an extraordinary improvement occurred in the physical signs. On the latter day the temperature suddenly fell to 97°; the ice was removed, and convalescence supervened. "It must be allowed, I think," it is stated, "that in this case the ice was of the greatest service. It is hardly too much to say that it saved the patient's life.'

CASE XLVI.—M., aged seven, had previously suffered from empyema, the right lung being entirely collapsed. The boy was first seen on the second day, when his temperature was 104°, and respiration 56. The pneumonia was believed to exist in the left lung, over which ice-bags were applied. The temperature fell four degrees before the next morning, but gradually rose somewhat during the subsequent forty-eight hours. On the third day the lips and cheeks were livid. This was not surprising, for the right lung was useless. On the fourth day the respiration was 60; on the fifth day the temperature was 102°, and respiration 58. On the seventh day the temperature fell to normal, and the ice-bag was removed.

CASE XLVII.—M., five years old, was admitted on the third day of his illness. The next day his temperature was 104°, and respiration 50. No signs of pneumonia could, however, be discovered until the sixth day, when it was found that the upper part of the right lung was consolidated. The ice-bag was applied until the eighth day, when crisis occurred.

CASE XLVIII.—M., about forty years old, had pleuro-pneumonia of the right base and pleurisy of the left side. On the second day the temperature was 104°, and respiration 32. The ice-bag was applied. Crisis occurred on the fifth day. This patient also had a very feeble heart. There followed a good recovery.

recovery.

CASE XLIX. (Reported by Mr. Back, Lancet, August 29, 1885).—M., thirty-five years of age, was seized with double basic pneumonia May 31, 1885; the temperature was 104°, the respiration 48. Hot flaxseed-meal poultices were applied, and 5 m, of tincture of aconite, fl.3ss of antimonial wine, and 5 m

of ipecacuanha wine were ordered every four hours, besides a diet of milk and beef-tea. On June 2d, the patient was much worse; the temperature was 106°, and the respiration 60; he was delirious, with profuse diarrhea, and was passing urine and feces involuntarily. On the 3d the diarrhea had ceased, but the man continued very delirious, and had a temperature of 106° and respiration 80. The whole chest was sponged with cold water every hour. On the 4th the temperature was 103°, the respiration 40, and the delirium was less. On the 5th the temperature was 100°, the respiration 34; the application of cold was continued. On the 6th the temperature was 99°, the respiration 30 From this time on the patient gradually convalesced.

CASE L. (Reported by Dr. F. Gundrum in The News, October 12, 1892, p. 473).—F., nineteen years old, suffered from pneumonia of the whole of left lung, with a temperature of 106° and a respiration of 46. On the first day the woman was wrapped, from chin to toes, in woollen blankets wrung out of cold water (50° F.), and changed every 30 minutes. At 1 0'clock the next morning the temperature was 102°; at 6 A. M., 98°, and respiration 16. She was now thoroughly rubbed, put in warm blankets, hot bottles placed about her feet, and half an ounce of brandy administered. This treatment aborted the disease completely.

The histories of these fifty cases open up many points of interest in the discussion of the influence of ice in the treatment of acute pneumonia, and as pertinent to this subject I will append the following comments:

1. The resolving power of ice on the exudation. This is a marked feature in its therapeutic action, and must be regarded as one of the strongest factors in its curative influence. This can at least be partly explained on the following basis: The most apparent lesion in croupous pneumonia is an enormous distention of the pulmonary capillaries, with partial or complete stasis of the blood in these vessels, exudation of the fluid constituents of the blood, and diapedesis of white and red blood-cells into the alveoli of the lung. It is well known that cold has the power of contracting the bloodvessels, and from this action one can understand why it should exert a beneficial action in pneumonia, by giving tone to the capillaries, by restoring the normal blood-flow, and thus checking the leakage. But there is often reason for believing that it also dissolves the exudate in the pulmonary alveoli. For example, there may be a pneumonic area in which there is absence of respiratory murmur, with the presence of a flat percussion note and bronchial breathing-indicating, beyond doubt, that the process has passed beyond the stage of engorgement and into that in which the exudation has taken place into the alveoli; yet the application of ice will, in a remarkably short time, develop a new group of physical signs, such as crepitation, reappearance of the respiratory murmur, diminution of flatness, etc., indicating that a break-down occurred in the exudation. This has not only been observed by myself, but is dwelt on by Dr. Lees, who says (p. 894): "In many ca es I noticed a striking arrest in the development of the physical signs," and that the ice-bag "distinctly tends to repress the inflammatory process in the lung."

Influence on symptoms. No less decided is the influence of the ice on some of the most prominent symptoms of pneumonia. The pain, difficult respiration, cough, and expectoration, are remarkably relieved, and the temperature is frequently depressed two or three degrees in the course of half a day. The beneficial influence exerted on these symptoms produces a very agreeable effect, and often makes the ice acceptable to those who, at first, protest against its use. This I have noticed in most of my cases, and it has also been witnessed by others, as will be seen by the histories of the cases that have been reported to me.

Is the ice injurious? My own rather limited experience with the ice-treatment does not show that it is accompanied or followed by any evil consequences, nor have any of those who have reported cases to me observed any such results, although some of them kept the ice applied for two weeks. Dr. Lees says (p. 894): "I have never seen any harm follow from the employment of the ice-bag in pneumonia."

Ages of patients. It is important to note in this collection that the ages of the patients in whose cases the ice was applied varied from infancy to old age—the youngest being six and a half months old, and the three oldest sixty, sixty-five, and seventy-four years respectively.

The results. It may be said, without claiming too much, that the results obtained from the icetreatment of pneumonia have been good. Out of the fifty cases which I collected but two were fatal, making a death-rate of 4 per cent. In estimating this mortality-rate it must be remembered that at least one of the cases (XIII) that died was an exceedingly unpromising one, being a sufferer from chronic lead-poisoning, and also very intemperate; while the pneumonia which caused the death of the other one (XIV) was in all probability an acute exacerbation of an old attack. In Dr. Lees' series of eighteen cases (from XXXI to XLVIII) no deaths occurred, nor did any occur in the eleven cases reported by Dr. Jackson. Moreover, the Lancet, of August 10, 1892 (p. 279), refers to an article by Dr. Fieandt, published in Duodecim, a Finnish medical journal (an original copy of which I am unable to secure), in which there is an account of 106 cases of pneumonia treated with ice-applications by that gentleman, and notwithstanding that among

these there were ten cases of double pneumonia, and that the epidemic of the disease was rather severe, he had only three deaths, or a death-rate of 2.82 per cent. Adding these cases to those reported in my collection, we have a total of 156 cases of pneumonia treated by means of cold applications to the chest, with five deaths, or a death-rate of 3 20 per cent.

While the number of cases reported here is not very large, it is nevertheless evident that the results of the ice-treatment are much superior to any other with which I am familiar. Thus, according to Osler, the mortality-rate of 1012 cases in the Montreal General Hospital was 20 per cent., while in the Charity Hospital at New Orleans it was 20.01 per cent. Of 1000 cases of pneumonia treated in the Massachusetts General Hospital, from 1822 to 1889, the mortality was 25 per cent. In Dr. Hartshorne's valuable paper on "Pneumonia" it is estimated that the death-rate from this disease in the Pennsylvania Hospital during the years 1884, 1885, and 1886, was a little more than 31 per cent. In comparing the results of the ice-treatment, so far as they go, with those which have been obtained from the treatment pursued in the hospitals named, we find that the former are about eight times better than the latter. It will be of great interest to see by future clinical investigation whether these satisfactory results can be maintained, and if this can be done even approximately, it is needless to say that a pronounced advance in the therapeutics of acute pneumonia has been made.

COUGH OF NASAL ORIGIN, WITH REPORT OF A

BY MAXIMILIAN HERZOG, M.D.,
LARYNGOLOGIST AND OTOLOGIST TO THE GERMAN HOSPITAL,
CINCINNATI, OHIO.

COUGH is, to a certain extent, a purely physiologic phenomenon, intended to support and supplement the action of the ciliated epithelial cells lining the respiratory tract. It is the function of these cells to keep up a current in the direction from the central organs of respiration—the lungs—toward the periphery, in order to remove the mucus secreted in the respiratory tract, and with it those mechanical adulterations, including pathogenic and non-pathogenic microörganisms, suspended in the surrounding air, that have gained access to the respiratory tract below the larynx by not having been effectually excluded by those organs especially equipped to do this, namely, the nasal cavities and the nasopharynx.

The act of coughing is considered physiologically as a modified expiration, brought about by reflex stimuli acting upon the pneumogastric nerve, more especially upon its branch, the superior laryngeal.

The pneumogastric nerve is to be looked upon as the guardian of the lower parts of the respiratory tract, and, as such, has to prevent detrimental elements from gaining access to the lungs, or to remove by reflex action those that have already gained access.

The act of coughing usually begins with a deep inspiration; then follows an expiration, which is, however, obstructed, in consequence of the approximation of the vocal bands as tightly as possible (more completely than in phonation), by the adductor muscles of the larynx. At the same time the abdominal muscles force up the contents of the abdominal cavity and the diaphragm, which itself contracts; and the air in the alveoli, bronchi, and trachea, which cannot escape through the tightlyclosed rima glottidis, is brought under a comparatively high pressure. Finally, the vocal bands relax, the air escapes forcibly, and its current, as it sweeps over the respiratory tract, carries along a certain amount of mucus and particles suspended in it, and, with the assistance of the muscles of the pharvnx. etc., delivers it to the outside world.

In perfect health, and when surrounded by pure air, cough plays an insignificant rôle, very much subordinate to such physiologic processes as, e. g., the act of micturition or that of evacuating the contents of the intestines. In many diseases, however, as, for instance, in croupous pneumonia, cough is the most important factor in ridding the alveoli and bronchioles of the pathologic products of disease. While a cough of this kind may be styled a necessary and beneficial reflex phenomenon, there are certain kinds of cough that are, from a physiologic as well as a pathologic point of view, of an absolutely useless character, dependent upon a reflex from a distant point, an irritation that, de facto, does not interfere, either sufficiently or not at all, with the act of respiration, and, therefore, cannot be favorably influenced by cough. Cases have been reported of cough of reflex origin from the liver and spleen (Naunyn), from the stomach and vesica urinæ (Krimer), from the external integument of the thorax (Leyden), from any and every part of the integument (Strubing), from the transmission of ordinary stimuli to the nerves of special sense (Ebstein)1.

During the last ten years it has been generally recognized that the nose is among those organs that may give rise to most obstinate cough of reflex origin. That pathologic changes in the nasal cavities constitute one of the most fruitful sourc s of various reflex phenomena is a fact that has been recognized for some time. The first, however, to mention this more elaborately and to lay especial stress upon it

¹ Quoted in Gottstein, Die Krankheiten des Kehlkopfes, 1850.

was Voltolini,1 whose views were afterward confirmed by reports of cases by B. Fraenkel, Hænisch, A. Hartmann, M. Schaeffer, Bresgen, J. Herzog, and especially Hack,2 who reported a number of instances and advanced a theory in explanation of reflexes of nasal origin. While the theory of Hack has been almost completely abandoned, he may still claim the merit that his paper was the most complete and most extensive on the subject up to his time. Among Hack's cases of reflex neuroses of nasal origin was also one of reflex cough dependent upon the presence of a nasal fibroid polypus. Carl Seiler³ followed with the report of two cases of reflex cough of nasal origin. John N. Mackenzie' next recorded his experiences on the subject. Thrasher5 reported two cases before the meeting of the Mississippi Valley Medical Association, at Louisville, in 1890. One case is also reported by Scheinmann,6 who tabulates and reports thirty-two cases of reflex neuroses of nasal origin occurring at the Throat and Nose Clinic of the University of Berlin, and whose extensive paper is undoubtedly the best contribution upon the subject which has appeared during recent years. Whether F. C. Heath, who contributes a paper on "Nasal Reflexes," in the American Lancet, Detroit, 1891, reports any new cases of reflex cough of nasal origin I am unable to state, as his paper has not been accessible to me.

Many of the authors who have written upon the subject of reflex neuroses of nasal origin point out the fact that it is almost universally in patients of a neurotic taint that these neuroses develop. Under some conditions the nasal cavities may show extensive pathologic changes; there may be new growths and points and surfaces compressing each other, without the appearance of any reflex phenomena. On the other hand, in individuals of anemic and neurasthenic type, slight changes, small areas of contact may bring about serious reflex disturbances. Some authors go as far as to maintain that no changes in the cavum narium of the kind usually producing reflex neuroses will do so unless developed upon the soil of an already disturbed nervous equilibrium.

If the normal nasal mucous membrane of an individual in perfect health, at the time not the subject of cough, and not accustomed to a prolonged course of nasal treatment, be gently irritated by a nasal sound, one will almost universally notice a twitching of the facial muscles of the side touched, a dilatation of the vessels of the nasal mucous membrane, and epiphora. If the irritation be more intense (such as may be caused by a weak electric current), sneezing, and, very often in nervous individuals, coughing will be excited. This observation is in favor of the claim of those who maintain that reflex neuroses of nasal origin only or almost universally develop on a neurotic soil. The case herewith reported may also be cited in support of this view:

M. B., an unmarried girl, twenty-one years old, of good family history, was said to have had "curvature of the spine" when she was four or five years old, for which she was treated, and got well. She suffered occasionally with pain in the back. For two or three years she had had a constant cough with a feeling of irritation and a sensation of the presence of a foreign body in the throat, of which she wanted to get rid. All treatment directed against the cough had failed to benefit her and ameliorate the constant irritation. The patient also stated that she sometimes had pain in swallowing, and had been told that she breathed through her mouth during sleep, though she did not snore. The sense of smell had always been good.

The girl was small and slender, looking less than her years. She limped slightly in walking, in consequence of a deformity of the spinal column in the lumbar region (probably a scoliosis—no direct examination of the spine having been made), and was

anemic, restless, and nervous.

The nasal mucous membrane was rather pale and anemic, and in a few discrete spots appeared atrophic. In the left nasal cavity, about three centimeters behind the anterior nares, close to the floor, was a mucous band, connecting the lower turbinated bone with the septum, stretching across and largely obliterating the inferior meatus. This band was from 3 mm. (at its middle part) to 5 mm. (at the two points of attachment) wide and of the same color as the neighboring parts. On the right side the mucous membrane of the lower turbinated bone showed some hypertrophy. The posterior extremity of this structure also showed a slight amount of thickening. Tenacious mucus covered the roof of the naso-pharynx. The pharyngeal mucous membrane was granular; on either side two granular bands extended from the plicæ salpingopharyngeæ downward; the band on the right side being especially well marked. Both tonsils were somewhat hypertrophic, the right one especially so. The epiglottis was imbedded in a hypertrophic lingual tonsil; the larynx was otherwise normal.

A diagnosis was made of adhesion in the left inferior nasal meatus, with chronic granular and lateral pharyngitis, and hypertrophy of the pharyn-

geal and lingual tonsils.

In view of the numerous hypertrophies found in the pharynx, it seemed justifiable to consider them as the cause of the cough, and they were consequently treated topically for a long time. The band of adhesion in the left nasal cavity was, on ac-

¹ Voltolini: Anwendung der Galvanokaustik, etc., Wien, 1871.

² Hack: Berl. klin. Wochenschr, 1882.

⁸ Archives of Laryngology, vol. iii.

⁴ Amer. Journ. Med. Sci., July, 1883.

⁵ Cincinnati Lancet-Clinic, October 11, 1890.

⁶ Scheinmann: "Zur Diagnose und Therapie der Nasen Reflexneurosen," Berl. klin. Wochenschr., 1889, pp. 295, 327, 399, 425, 471.

count of its situation and color, readily overlooked, and was only discovered when the patient came to my clinic. Touching the mucous band connecting the lower turbinated bone with the septum did not excite a fit of coughing. This observation did not, however, at all influence me in my belief that in the adhesion I had found the cause of the reflex cough. It has frequently been noticed that pathologic changes in the nose act as the cause of reflex neuroses that disappear after proper surgical interference or other therapeutic measures, even if, on touching these structures or tender points, as the case may be, the reflex is not always excited. The result of such interference in the case reported in this paper proved the correctness of the assumption.

Two days after the examination the lower parts of the left nasal cavity were treated with a ten per cent. solution of cocaine, and the band of adhesion, which now appeared very much on the stretch, was divided by a galvano-cautery platinum knife. No pain was experienced, and no hemorrhage followed. Insufflation of europhen was practised after the

operation.

On the following day the patient reported that the cough had disappeared completely. The good effect continued. A ferruginous tonic was prescribed, but the pharynx was not touched.

The nasal mucous membrane at the points of cauterization healed kindly; the cough did not

reappear.

The patient, after taking iron for some time, felt better than she had for years. She looked well, had gained in weight, and her cheeks presented a rosy color. The hypertrophies of the pharynx still persisted, as they had not been interfered with surgically. The cough, which had left the patient directly after the operation, had not reappeared.

The case is instructive not only as one of cough of nasal origin, but also because it shows that hypertrophies in the pharynx, which often give rise to reflex symptoms, may not do so, even in a nervous subject in whom structural changes (in our case an obstructive adhesion) in the nose will do so.

In conclusion, I would like to state that it is my belief that the adhesive band in this case originated from ordinary hypertrophies, such as are found in almost every case of rhinitis hypertrophica. Probably a point of contact was established between the mucous membrane of the turbinated bone and that of the septum, a slight ulceration occurred in consequence of the hypersecretion dependent upon the hypertrophy, and the adhesion was formed at the point where two ulcerated surfaces were in contact. Later on the hypertrophy underwent a retrograde change (there being, already, spots of a somewhat atrophic character noticeable at the time the patient presented herself), and the mucous membrane contracted, but left behind a band of adhesion at the place that formerly had been a point of contact, the band acting as a source of reflex cough.

No. 50 W. NINTH STREET.

CLINICAL MEMORANDA.

TRAUMATIC EMPYEMA COMMUNICATING WITH THE BOWEL.

BY N. T. DULANEY, M.D., OF BRISTOL, TENN.

E. G., thirty years old, a conductor on a freight train, while engaged in the discharge of his duties was shot with a 38-caliber pistol at short range, the muzzle of the pistol being within a few inches of his body. The ball entered his body in front, at a point about five inches below the left nipple, and one inch and a half outside the left nipple line. The ball could not be located for several weeks. I saw the patient nearly a month after the reception of the injury, in consultation with his attending physician, Dr. M. M. Butler, of Bristol, Tenn. I learned that the patient had not had very much fever, the temperature varying from 99° to 101°; the pulse from 90 to 100, and not sthenic. When I saw the man the temperature was 102°. There was very little pain; no cough, nor had there been any at any time; no bloody expectoration to indicate a wound of the lungs; little or no appetite. The bowels were torpid: the breathing somewhat hurried and more pronounced on the right side. The patient was considerably emaciated.

Physical examination revealed extensive dulness over a large portion of the left chest, amounting to flatness, with absence of normal respiratory sounds, and increased percussion-resonance and exaggerated respiratory murmur on the right side. The case not improving, but symptoms of septicemia becoming more pronounced, I was again called twelve days later. Physical examination disclosed the same condition as before, but somewhat aggravated and with the addition of a distinct prominence as large in circumference as the mouth of an ordinary teacup, situated on the back, to the left of the spine, and corresponding to the eighth and tenth ribs in this region. With this as a guide, the patient was anesthetized and a vertical incision about three inches in length made through the prominence, down to the ribs. There was now a gush of very offensive, dark fluid, amounting to about a quart, which seemed to be a mixture of serum, pus, and the débris of disintegrating tissue.

The ball was found just outside the chest under the deep muscles of the back, in its passage having broken a rib-the tenth, I think-about one and a half inches from the spine. A portion of the same rib was broken, but not detached, for a distance of about an inch, and was hanging loosely in the chest in such a way as to obstruct the opening in the chest-wall. This being removed with forceps, the opening was at once filled by mass of dead tissue forcing its way out. I began drawing this out and continued until no more came, when I had as much as I could hold in my two hands. It was a stinking mass of dead tissue, consisting, I think, of pleura and lung and connective tissue. It was dark, almost black, thickly covered with grayish, ash-colored spots as large as a wheat grain and larger. Having removed all of this dead tissue, the cavity was washed

¹ Read before the Tennessee State Medical Society, April 11, 1893.

out with a weak, warm solution of carbolic acid, and the wound dressed with iodoform-gauze, absorbent cotton, and bandage, and the patient quieted with a dose of morphine hypodermatically. The temperature steadily declined, the wound being dressed every day and the cavity washed out with warm water, usually with a little carbolic acid added.

The day after the operation we used an enema to evacuate the bowels. The nurse, a negro man, said that the water of the injection passed out through the opening in the back. This being doubted, the physician in charge tested it the next day by injecting into the bowel water colored with potassium permanganate, and he found that the fluid injected into the rectum passed freely and promptly out through the opening in the back. This was an unexpected and perplexing complication.

From this time on the wound was dressed and washed out every day for some weeks, and for three or four months fecal matter passed out through the wound freely, especially when the discharges were thin,

Two months after the accident I dressed the wound, and a mass of fecal matter as large as a hulled walnut passed out. The patient slowly but steadily improved, and is to-day on duty as a railroad conductor, looking well and weighing 193 pounds, the fistula having entirely closed.

This is an interesting case, because:

1. There were no rational signs pointing to injury of the lung. There was no blood expectorated, no cough, and at no time was the sputa characteristic of pneumonia. The case went on for seven weeks, the chest filling with serum and pus, and a large amount of tissue undergoing necrosis within the chest. Yet, aside from the physical signs and the altered breathing, there was nothing to indicate the exact nature of the case. It is true, the persistent fever, emaciation, loss of appetite, restlessness, and failing strength indicated something seriously wrong, but it remained for the physical examination to determine the nature of the condition.

2. The ball in its passage had so damaged the diaphragm and the bowel at some point in contact with it as to lead to inflammatory adhesion, followed by sloughing, and resulting in a fecal fistula. With all this, there was no bloody or mucous discharge from the bowels, and no symptoms indicating injury of the bowel until the water injected into the rectum was seen to pass out through the wound in the back.

In regard to the management of this case, while no new principle is involved, the result forcibly emphasizes the two or three main points in the local treatment of such cases, whether of idiopathic or of traumatic origin—namely, cleanliness, or antisepsis, if you please, with thorough drainage. These given, unless your patient is exhausted from protracted disease, or septicemia, or some undermining constitutional disease, you are in most cases warranted in making a favorable prognosis.

It is astonishing how promptly convalescence sets in after the inauguration of such treatment, if not too long delayed, or if the ordinary vis medicatrix natura is present to assist you. Of course, constitutional treatment, based on general principles, is not to be neglected.

The distance from the point of entrance of the ball to the point where it was cut out around the chest is eight and three-quarters inches.

CYSTIC DEGENERATION OF ENLARGED SURMAXILLARY GLANDS.

BY D. W. HAYS, M.D., OF OSAWATOMIE, KAN.

MRS. B., born in 1819, when about thirty years old noticed a "kernel" under the right jaw, which, however, did not subside in the usual manner, but continued to slowly enlarge, though it remained freely movable. It gave no inconvenience or pain, and only after it had attained considerable size was anything done to check its growth or produce its absorption. Then, only local applications of tincture of iodin and the like were used until 1871, when the mass had become a little larger than a man's fist. A diagnosis of multilocular cyst was made, and removal advised. The opinion was expressed that the growth would not undergo further enlargement, or shorten the woman's life, and she therefore refused operative interference. Contrary to prediction, the mass continued to increase in size, and even more rapidly than before, so that at the time of death, Nov. 10, 1802, at the age of seventy-three years, it measured fifty inches in circumference and, after removal, weighed forty-seven nounds.



The mass was easily removed, being encapsulated and attached only to the integument and connective tissue. The superficial veins were numerous and quite large; the arterial supply was small, and the vessels were not traced to their origin. One large branch from the facial nerve, with numerous dividing branches, was found on the upper part; other nerve-supply was not sought. The inferior maxillary bone had been pushed or dragged forward, and had been absorbed from below until it presented a decided upward curve.

The contents of the tumor were found to be a greenishyellow fluid. The walls were, at some portions, thinned until they were little more than half an inch thick; at no place were they more than an inch and a half thick. The tissues of the walls seemed to be thoroughly disintegrated and of a cheesy consistency. A section from the apparently best organized portion showed nothing distinctive under the microscope, except some broken-down glandular tissue.

The accompanying photograph was taken some eight or ten years ago, and does not show the actual size at death; it would have shown better if it had been suspended.

I think this case worthy of record on account of the large size and situation of the growth, and especially because of its evident origin, as it is quite certain, from the best history that I can obtain of its commencement, that it was an enlargement of one of the submaxillary glands that became cystic by degeneration,

MEDICAL PROGRESS.

A Case of Diphtheric Croup in which a Tracheotomy-tube was Worn for Sixty Days.—At a recent meeting of the Philadelphia Academy of Surgery, DR. H. R. WHARTON reported the case of a child eighteen months of age, with diphtheria, in which an intubation-tube was introduced for the relief of dyspnea. At the end of fifteen hours the instrument was coughed up and the dyspnea returned. Tracheotomy was then performed, with relief to the dyspnea, and the patient subsequently did well.

On the tenth day after the operation the tracheotomytube was removed and the patient breathed comfortably for a time, but after fifteen or twenty minutes the dyspnea recurred and became so urgent that the tracheotomy-tube had to be replaced.

Attempts were made to remove the tube at intervals of a day or two for a week or more, with a similar result. Four weeks after the operation the child was etherized and the tracheotomy-wound was dilated, so as to expose the wound in the trachea. It was then found that there were a number of masses of granulation-tissue springing from the trachea in the region of the tracheal incision; these were removed with forceps and scissors, and the tracheotomy-tube was again introduced. After waiting a few days another attempt was made to remove the tracheotomy-tube, but this also failed. An intubation-tube was introduced upon the removal of the tracheotomy-tube, and was worn for some hours, and upon its removal the child was able to breathe comfortably for five hours; but after this time dyspnea recurred, and the tracheotomy-tube was again introduced. After several trials of the intubation-tube, it was finally abandoned. The patient was again etherized, the tracheotomy-wound enlarged, and a number of masses of granulation-tissue removed, and their bases were touched with the solid stick of silver nitrate. An intubation-tube 'was then introduced, and was worn for twelve hours, when it was coughed up. Thinking its expulsion was probably due to the irritable condition of the larynx and trachea, resulting from the recent operation, the tube was not replaced for forty-eight hours. To keep the tracheotomy-wound open, so that the tracheotomy-tube could be introduced, if it were required, an obturator was introduced into the tracheotomy-wound. The intubationtube was worn for four days and was then removed, and the obturator was retained in the tracheotomy-wound for

three days longer, and was then removed. The child after this time had no further difficulty in breathing, and the tracheotomy-wound, after the removal of the obturator, rapidly contracted and healed. The patient wore the tracheotomy-tube for a period of sixty days.

Rupture of the Uterus during Labor, with the Placenta in the Abdominal Cavity .- WASTEN (St. Petersburger medicin, Wochenschrift, No. 19, 1893, p. 173) has reported the case of a woman, twenty-nine years old, who was delivered of her ninth child with the aid of forceps. after a not unduly protracted or difficult labor. The child was dead. The placenta had not been expelled. Examination showed the uterus to be contracted, but the placenta was in the abdominal cavity. Notwithstanding the precarious condition in which the woman was, celiotomy was undertaken. Cocaine was injected in the line of the contemplated incision and chloroform was cautiously administered. The portion of the umbilical cord in the vagina was cut away, and the placenta, with numerous blood-clots, was removed from the abdominal cavity. Then the uterus was amputated above the vaginal portion and the broad ligaments and large vessels ligated. The stump was gently cauterized with the hot iron and replaced in the abdominal cavity. Drainage ber vaginam was provided for and the wound closed. Convalescence was attended with no noteworthy complication, other than the occurrence of an endometritis in the remaining portion of the cervical canal and the formation of an exudate in the right iliac fossa. It was found that the rupture had involved the anterior wall of the uterus in the situation of the placental insertion. It is thought to have taken place spontaneously.

Qualitative Variations in the Knee-jerk,-BENEDIKT (Neurologisches Centralblatt; Deutsche medicin, Wochenschr., No. 19, p. 464) describes four varieties that the knee-jerk may present: (1) It may be clonic; several jerks rapidly succeeding one another after a single tap upon the tendon. This may occur in cases of palsy, of myelitic or cerebral origin, and with spastic conditions of spinal origin. (2) The knee-jerk may be paradoxic, the leg being jerked in flexion instead of in extension. This has been observed in a case of tumor of the roof of the fourth ventricle at the level of the striæ acusticæ, presenting headache, static vertigo, and unilateral nervous deafness, to which melancholic apathy was subsequently added. A sub-variety of this form consists in the occurrence of a flexor jerk following the extensor jerk. (3) Percussion of the quadriceps tendon is in some cases followed by contraction not only of the extensors or flexors of the knee, but also of some of the muscles of the trunk and of the opposite side of the body. (4) The knee-jerk may be tonic; the leg is slowly extended, but fails to fall back to its original position. There is, however, no actual spasm of the quadriceps. In cases of marked lateral sclerosis, with greatly increased reflex irritability, percussion of the quadriceps may induce tonic extensor or flexor spasm. (5) The knee-jerk may be delayed; or (6) it may become exhausted.

Primary Sciatica.—GIBSON (Lancet, No. 3633, p. 860) has reported the results of an analysis of 1000 cases of

primary sciatica, with especial reference to the treatment of 100 cases by acupuncture. Of the whole number. 884 cases occurred in males: 116 in females. In 443 the right side, in 483 the left side, and in 74 both sides were affected. In 14 cases the affection was first observed between the ages of 15 and 20; in 150, between 21 and 30; in 310, between 31 and 40; in 248, between 41 and 50; in 187, between 51 and 60; in 71, between 61 and 70; in 11, between 71 and 80. In 132 there was accompanying lumbago. Of 100 consecutive cases of sciatica treated by accouncture, 56 were cured, 32 were much improved; 10 were improved; and 2 were not improved. A single spear-pointed needle two and a half inches long, was employed. This was plunged into the nerve itself, but immediately withdrawn. The nerve should be pierced about five times over the parts where there is pain on pressure: this can be accomplished through one cutaneous puncture by partially withdrawing the needle and entering it again at a somewhat different angle. A single puncture may suffice. The measure may, however, be repeated at intervals of a few days. Rest is an essential factor in the treatment. A gouty or a rheumatic diathesis should receive appropriate attention.

Renal Hematuria, Lasting Twenty Years .- WETMORE (Maritime Medical News, vol. v, No. 5, p. 65) has reported the case of a man, sixty-four years old, who for twenty years, intermittently, passed blood in the urine. The first attack occurred in the winter, without known cause, and lasted for two months. The second attack occurred three or four years later, and also commenced in the winter and lasted for two months. The attacks gradually became more frequent and lasted longer, the urine for six years not having been free from blood for more than a few weeks at a time. There was no family history of carcinoma or tuberculosis, and no personal history of malaria. There had never been colic. In some attacks there were pain and tenderness in the right loin and pain and swelling in the right hip and lower extremity, and swelling of the left lower extremity. The urine contained a moderate amount of albumin and many red blood-corpuscles and some blood-casts. An ultimate diagnosis was not arrived at,

Acute Pulmonary Hypertrophic Osteoarthropathy .- At a meeting of the Société Médicale des Hôpitaux, Moi-NARD (La Semaine Médicale, No. 13, 1893, p. 247) reported the case of a child, six years old, which three months after an attack of meta-pneumonic purulent pleurisy, presented deformity of the terminal phalanges, characterized by lateral thickening of the extremities of the fingers, which looked like drum-sticks. The nails were curved longitudinally and transversely like a parrot's beak or a watch-glass. They were abnormally deep red in color and extremely friable, but not striated. There was no other bony lesion. A second case occurred in a girl of five. The sequence and the lesions were quite like those in the first case, and the deformity disappeared with the cure of the pleuro-pulmonary condition.

MARFAN related that he had observed three cases of similar kind; one of empyema, one of bronchiectasis, and one of pyelo-nephritis.

THERAPEUTIC NOTES.

The Treatment of Suppurating Tuberculous Tissues .- A new treatment for suppurating tuberculous tissues has lately been suggested by JEANNEL, of Toulouse. The diseased tissues are freely exposed and, after the surrounding healthy parts are protected by a damp towel. the diseased structures are thoroughly sponged with a tampon of hydrated cotton-wool previously placed in a boiling saline solution. The application should be made to all irregular surfaces. After the procedure has been repeated five or six times it may be concluded that all of the tubercle-bacilli present have been brought into contact with the boiling solution, and have thus been destroyed. It is stated that this method has been employed in some cases in which amputation seemed to be the only remaining resource, with a satisfactory result and without mutilation .- Medical Press and Circular. No. 2820, p. 547.

Bromoform-intoxication .- DEAN (Lancet, No. 3636, p. 1062) has reported the case of a girl, four years old, with whooping cough, for whom was prescribed a four-ounce mixture, containing sixty minims of bromoform, and of which it was directed that two drams should be taken every three hours. In the course of the treatment the physician was hastily called to see the child and found it with pin-point pupils, livid, and breathing somewhat stertorously. On inquiry it was learned that the attendants had failed to thoroughly shake the mixture before its administration, and that, as a consequence, the child had received with the last dose from the bottle about fifteen or twenty minims of bromoform. The stomach was evacuated by means of a pump, the contents smelling strongly of bromoform. Strong coffee was injected into the bowel, and a small dose of apomorphine injected beneath the skin, and the battery employed. The child slowly recovered and was shortly quite well.

For Tuberculosis.—

R.—Iodoformi gr. xx-xl.

Essentiæ eucalypti
Essentiæ terebinthinæ
Essentiæ guaiaci
Creasoti

S.—By inhalation or spray; or from five drops upward Petresco, La Roumaine Méd.; Münch. med. Woch.

Calomel for Gout.—GRIMM (Deutsche medicin, Wochenschr., No. 17, 18; Wiener medicin. Presse, 1893, No. 2, p. 818) has successfully employed calomel in a single large dose, in the treatment of acute and subacute gout. The result is ascribed to the stimulation of peristaltic activity, as it is known that an attack of gout is often preceded by intestinal torpor. Yellow mercuric iodid was found to act even more energetically than calomel.

To Allay the Craving for Alcohol.-

R.—Tinct, capsici f 3ij.

Tinct, nucis vomicæ
Acid, nitrohydrochlor, dil.

Infus. gentian. . . ad f3xij.—M. S.—Two tablespoonfuls as required.

The Practitioner.

THE MEDICAL NEWS.

A WEEKLY JOURNAL
OF MEDICAL SCIENCE.

OMMUNICATIONS are invited from all parts of the world.
Original articles contributed exclusively to THE MEDICAL NEWS will upon publication be liberally paid for,
or 250 reprints will be furnished instead of payment. When
necessary to elucidate the text, illustrations will be provided
without cost to the author.

Address the Editor: GEO. M. GOULD, M.D.,

PHILADELPHIA.

Subscription Price, including Postage in North America.

PER ANNUM, IN ADVANCE \$4.00.
SINGLE COPIES 10 CENTS.

Subscriptions may begin at any date. The safest mode of remittance is by bank check or postal money order, drawn to the order of the undersigned. When neither is accessible, remittances may be made, at the risk of the publishers, by forwarding in registered letters.

Address,

LEA BROTHERS & CO., Nos. 706 & 708 SANSOM STREET,

PHILADELPHIA.

SATURDAY, JUNE 24, 1893.

THE REPORT OF THE COMMITTEE ON REVISION OF THE CODE.

THE MEDICAL NEWS cannot agree with either the arguments or the recommendations of the majority of the Committee on Revision of the Code of the American Medical Association, in so far as these apply to the regulations governing the conduct of physicians. We have already stated our belief that the paragraphs referring to the conduct and obligations of others than physicians have no proper place in the Code; and in this we are at one with the majority of the Committee. When, however, that majority recommends that physicians should be permitted to patent mechanical devices, and to consult with all "legalized practitioners," we must emphatically dissent. On these two propositions the coming battle is to be fought; and the lines should be distinctly drawn.

As to the patenting of instruments, the objections are many; and the alleged parallel with copyrighting of books does not hold. The relation of an author to his readers is that of teacher to pupil. He devotes much time and study to research, literary and otherwise, and expresses the results in a certain manner. For this he receives a compensation, often quite inadequate to the labor performed. [We are speaking now, of course, only of the best books;

not of the books written with paste-pot and shears.] That compensation is exactly parallel with the fee paid lecturers and demonstrators by their pupils. It differs from a royalty on a patent in that it is not paid for permission to do certain things, but for instruction how to do them. For example, one might pay a fee for instruction in the performance of intubation; but this would be very different from a royalty on the instruments used.

The copyright has a further most important aspect, as security to the publisher that, having expended certain sums and incurred large risks in the preparation of a book, his investment shall not, in case the book proves successful, be stolen by others who have taken no risk. Beyond this, too, it protects an author from misrepresentation through careless book-making or unauthorized omission or addition; and thus protects the student from being misled.

Patents, on the other hand, are in the nature of tribute. They have invariably a tendency to unduly increase the cost of an instrument, to restrict its application, and to prevent its improvement. The only thing to be urged in their favor is the increased revenue to the owner of the patent. No better illustration of the purely commercial purpose of patents can be given than the history of the "pneumatic cabinet," for which at first a yearly rental was demanded equal to the sum for which the apparatus was afterward sold outright when its claims were found to be exaggerated.

If the sole purpose of the medical practitioner is to make all the money possible out of medicine, then let him patent his instruments, as any other tradesman would. Let him also patent his prescriptions if possible; for there is no essential difference, moral or intellectual, between altering the curve of a knife-blade and altering the proportions of calomel and jalap in a powder. The true parallel is that between patented medicines and patented instruments; not between patented instruments and copyrighted books.

Should the wise restriction of the Code be abrogated, what a delightful time we can expect when Professor Biggun hauls Doctor Greathead into court for infringing the patent of his obstetric forceps or his nasal cotton-holder. The "Priorität-streiten" of our Continental colleagues will sink into insignificance. And what a field for expert testimony! If Professor Biggun's students should swear that he lectured on the instrument in 1880, who knows but

that Doctor Greathead will exhibit a picture of it in an Italian tome of 1654?

Supposing the physicians of America to be anxious to lose all dignity and self-respect in a mad scramble for the Almighty Dollar, certainly there is but one means more effectual than the patenting of mechanical devices, and that is offered by the proposed provision as to consultations.

The special plea of the majority of the Committee that to consult with an irregular does not mean to hold a regular consultation equals in logic and humor the old conundrum which declares a door not to be a door when it is aiar.

We are not opposed to specialists or specialism; we believe in both; but we are opposed to that narrow view which places the specialist as a god upon a pedestal of superior wisdom, and makes the family physician but a nurse, or at best a clinical clerk who rehearses a history. No specialist worthy the name but consults with the attending physician, actually as well as nominally. If any refuses to do this, he is an excellent man not to send patients to, The attending physician is the one responsible for the management of the case, whether his consultation be in a case of cataract, or a case of ovarian tumor, or a case of enteric fever. He selects the consultant in each instance for the latter's special knowledge and experience in that particular class of cases. And whether the outcome of the consultation be an operation by the surgeon or the bathing of the patient by the nurse, in each instance the one who performs the work does so as the agent of the attending physician and by his consent and direction.

Such being the case, no loophole of escape from the full meaning of consultation can be found in the case of a specialist treating the patient of an irregular practitioner in professional association with the latter. The irregular is the responsible attendant, the specialist is his agent in performing the necessary manipulations. There is no other possible view. We have on other occasions shown why this or any professional relation with quacks or sectarians is unwise and immoral. While for the present merely referring to our former utterances, we may at some future time recur to the subject, for the reason that instruction upon the issues involved seems to be needed.

We trust that those interested in the preservation of professional purity will act on the advice of PRESIDENT MCGUIRE and secure discussion and for-

mal expressions of opinion in their respective County and State societies, in ample time to influence in the right direction the action of the American Medical Association at its next meeting. Those who wish the door ajar for unrestricted consultations will certainly be active and wary; those opposed to this degradation should be no less so.

NEPHRITIS OF MALARIAL ORIGIN.

As our knowledge of disease-causes grows, our perception of disease-complications correspondingly enlarges. It is thus that the number of so-called idiopathic affections becomes constantly less.

It is only since the studies of LAVERAN have incontestably shown that malarial disease is dependent upon the presence of parasites in the blood, and since we have by direct observation and by analogy become familiar with the biology of these minute organisms, that we have learned to recognize and understand the complications and sequelæ of an important group of diseases. While this knowledge has added nothing to our therapeutic resources, it has placed upon a rational basis a method of treatment that, however successful, has hitherto been purely empiric.

We are now fully able to appreciate why anemia should be so common a manifestation in the course of malaria; why under certain conditions of virulent intensity the pernicious forms should terminate fatally; why under other conditions thrombi should form in the bloodvessels and give rise to symptoms as variable as the organs in which the morbid process takes place. By the gradual accretion of knowledge of this kind we are prepared for the announcement that malarial infection may give rise to nephritis. Of this interesting association three instances are reported by STEFANOWICZ (Wiener klin. Wochenschr., No. 20, p. 365). In each the subject had lived in a malarious region; specific organisms were found in the blood; all of the symptoms, including the presence in the urine of albumin and casts, disappeared upon the administration of quinine in antiperiodic doses.

Three explanations of these remote or secondary complications of malaria at once suggest themselves: First, a toxic process; second, thrombosis; third, hemorrhage. Without here entering into a discussion of the various possibilities in the case, we venture to express the view that the first proposition is that which in most cases best explains the conditions present. There is much reason to believe

that in diseases of malarial origin toxic products are set free in the circulation, primarily as a result of the activity of the parasites present, and secondarily as a result of their destructive influence upon the red blood-corpuscles.

The subject is one of practical interest, and worthy of further study. In this connection it is interesting to note that methylene-blue has already been recommended in the treatment of nephritis as well as of malarial disease.

EDITORIAL COMMENTS.

Stretching of the Facial Nerve for the Relief of Trigeminal Neuralgia.-Among the most obstinate of intractable affections are the neuralgias, and of these perhaps the most intractable is that of trigeminal type. At the recent German Surgical Congress, SCHULZE-BERGE (Deutsche medicin. Wochenschr., 1893, No. 22, p. 535) detailed a mode of treatment which, despite the admonition of some distinguished men present, still appears to have much to commend it, and to be deserving of trial before resort to major measures. The procedure consists in stretching the facial nerve of the affected side, in the hope of removing a source of irritation dependent upon the functional activity of this nerve. This practice was carried out in the case of a woman who had suffered for six years from neuralgia of the third branch of the trigeminal nerve, and in which extirpation of the buccal nerve had been unattended with relief. The pain ceased after the fifth day. In the subsequent discussion, Esmarch and Gussenbauer maintained that the stretching should not be performed until other measures, including a course of castor oil, had failed. König pointed out the possibility of permanent paresis as a result of the operation.

The Etiology of Friedreich's Ataxia. - While there is practical unanimity of view as to the clinical history of the affection described by Friedreich and known by his name, or in consequence of deficient knowledge, as hereditary ataxia, there is some diversity of opinion as to the pathologic basis of the disorder. The prevalent view is that the symptoms depend upon sclerosis of the posterior and lateral columns of the spinal cord, though the factor of heredity is probably as often absent as present. SENATOR (Berliner klin. Wochenschr., 1893, No. 21, p. 489) reports a typical case, but dissents from the current view of its pathology. He presents facts and advances argument to show that the symptoms are to be referred to a congenital defect of development of the cerebellum, to which the changes in the posterior and lateral columns may be secondarily added, but are not essential.

Professional Confidence.—A propos of our recent remarks on the sanctity of the confidence reposed by patients in physicians (THE NEWS, April 22, 1893, p. 439), it is interesting to note that a bill has been introduced into the Canadian Legislature to place the physician in this connection in the same category as the

lawyer, and to regard as privileged confidential communications between patient and physician.

Similar legislation, it appears, has already been enacted in New York and in Michigan. In France it is penal for a priest, lawyer, physician, druggist, or midwife to reveal information into possession of which he or she may have come in the course of professional duty.

Bacillus Aerogenes Meningitidis.—In the cases of cerebrospinal meningitis hitherto studied from a bacteriologic point of view, the pneumonia-coccus (micrococcus lanceolatus) is the organism that has been most frequently found present. In two cases recently studied, an Italian observer, Centanni, has found exclusively an organism to which he gives the name bacillus aërogenes meningitidis.

Ecphyaditis is the name proposed by Dr. Robert T. Morris as a substitute for appendicitis. The word is a good one, as the term ecphyas already exists in medicine to describe the appendix vermiformis ceci. Ecphyaditis is also philologically the preferable word; it has the further advantage of distinctiveness.

REVIEWS.

THE TWELVE TISSUE REMEDIES OF SCHÜSSLER, COM-PRISING THE THEORY, THERAPEUTIC APPLICATION, MATERIA MEDICA, AND A COMPLETE REPERTORY OF THESE REMEDIES, HOMEOPATHICALLY AND BIO-CHEMICALLY CONSIDERED. By WILLIAM BOERICKE, M.D., late Professor of Materia Medica in the Hahnemann Hospital College of San Francisco, etc.; and WILLIS A. DEWEY, M.D., Professor of Materia Medica in the Hahnemann Hospital College of San Francisco, etc., 3d edition. Rewritten and enlarged. 8vo, pp. 384. Philadelphia: Boericke & Tafel, 1893.

THE remarkable confusion of terms exhibited in the title of this volume is an index of the confusion of thought that pervades it. It is possible for one to honestly believe the dogmas of homeopathy; but not to believe those dogmas and at the same time to think clearly. It seems that one Dr. Schüssler, of Oldenberg, Germany, in 1873, published an article entitled: "A Shortened Homeopathic Therapeutics," in which he undertook to prove that all diseases curable at all were curable by the administration in infinitesimal doses of those mineral substances that form part of the normal animal body. The "Twelve Tissue Remedies" are "ferr. phos., calc. phos., natr. phos., kali phos., kali mur., natr. mur., calc. fluor., silicia, calc. sulph., natr. sulph., kali sulph., magnes. phos." By a table published on page 29, we see that these salts are constituents of various other drugs, such as "china, gelsem., berb. vulg., hamam., cimicif., coloc.," etc., etc. From all of which, by pure Hahnemannian logic, it follows that Schüssler is right in his practice, but wrong in claiming, as he now does, that he has gone beyond homeopathy, and founded a new "new school" of Bio-chemiopathy-or something of the sort. There are several practical wheatkernels in this barnful of chaff, but the result is not worth the labor of winnowing.

LECTURES ON MENTAL DISEASES, DESIGNED ESPECIALLY FOR MEDICAL STUDENTS AND GENERAL PRACTITIONERS. By HENRY PUTNAM STEARNS, A.M., M.D., Physician Superintendent of the Hartford Retreat, Lecturer on Mental Diseases in Yale University, etc. With illustrations. Pp. 627. Philadelphia: P. Blakiston, Son & Co., 1893.

THIS volume consists of twenty-nine lectures, which the author delivered before the students of Yale University, and covers the field of mental diseases. The tone of the lectures is conservative, representing, for the most part, the consensus of opinion of American alienists. The classification of insanity is not so extensive as that adopted by Spitzka and other modern writers. The existence of paranoia as a special form of insanity is vigorously combated by the author, while such other special forms as katatonia, confusional and stuporous insanity are not even mentioned. We are glad to note that the author advocates the treatment of some cases of insanity outside of the asylum.

Some eighty pages at the end of the book are devoted to extracts from the laws of the different States and Territories of the United States which relate to the general care of the insane. Unfortunately, these are, in some instances, not given at sufficient length to be of much value.

Taken all in all, the book is of such a character that it would enable a student or general practitioner to obtain a good working knowledge of insanity and would supply him a safe guide for treatment. The specialist will find nothing new in it save the notes of the author's experience at the Hartford Retreat, of which some of the cases recorded constitute a real contribution to the science of psychiatry.

THE ANATOMY OF THE PERITONEUM. By FRANKLIN DEXTER, M.D., Assistant Demonstrator of Anatomy, College of Physicians and Surgeons, New York. 12mo, pp. 86, with 38 illustrations. D. Appleton & Co., 1892.

This is a valuable contribution, intended to render more clear that unsatisfactory and incomprehensible anatomic part, the peritoneum.

The author, taking the description of Prof. Toldt, of Vienna, of the development of the peritoneum, as a basis, has enlarged the work with many plates, and has so planned it that the student is taken along, step by step, in the most elementary manner, with comparative ease, to a perfect understanding of the subject. It is not sought to impress the embryologic details, but their results. The study of the development of the organs not only contributes valuable information as to their normal position, but renders the knowledge of abnormalities more comprehensible. Thus, the hiding from view of a portion of the duodenum by the mesentery of the transverse colon, the absence of peritoneum on the posterior surface of the mesentery and a portion of the colon, the high position of the cecum in children in relation to the liver rather than to the iliac fossa, the simultaneous development of the peritoneum and the intestine (that the intestine has not, after it has been formed, been pushed into the membrane as a finger into a glove), and many other points are satisfactorily explained.

The work is of a highly scientific character and of considerable practical value, and the author is to be highly commended for the production of a work which reflects great credit upon his ability as a practical scientific anatomist.

PHYSIOLOGY. A MANUAL FOR STUDENTS AND PRAC-TITIONERS. By FREDERICK A. MANNING, M.D. (The Students' Quiz Series.) Philadelphia: Lea Brothers & Co.

This little book claims to be but a brief summary of the salient features of human physiology. This is the statement with which the writer begins his preface, and it requires but slight acquaintance with the book to prove that he appreciates the proper sphere of the quiz compend. The terse answers to questions, and the descriptions which he gives, while they contain the pith of the subject, could not be made to take the place of the study of one of the standard text-books; but once the information is acquired from such a source, this manual will be an excellent help to fix it in the mind of the student preparing for his examination, or, perhaps, to refresh the memory in after years.

On looking through the various sections of the book—on the Blood, the Circulation, the Respiration, Digestion, Absorption, Animal Heat, Secretion, Muscle, Nutrition, the Nervous System, the Senses, and Embryology—we confess to be well pleased with it, as a whole. Of course, everything is much condensed, as it is proper that it should be, but the author has succeeded very well in presenting the essence of the subject. In an appendix are given a Table of the Development of the Embryo, a list of the Chemical Tests used commonly in Physiological Analysis, and the Metric System, which are well to have at hand.

Dr. Manning has based his book mainly on the works of Dalton, Foster, and Kirke. He has written in a very readable style, and deserves to be congratulated on his contribution to the Quiz Series.

TRANSACTIONS OF THE AMERICAN GYNECOLOGICAL SOCIETY. Vol. xvii, 1892. Philadelphia: William J. Dornan, 1893.

THIS volume presents in the usual complete form several interesting contributions, among which are to be noticed particularly the extensive paper upon "Tuberculosis of the Female Generative Organs," by J. Whitridge Williams, of Baltimore, Md. This article is very extensive in its scope and presents a complete history of the investigations in this field of Gynecology. The writer naturally devotes most of his space to the consideration of tuberculosis of the tubes, as these are the primary seat of tuberculous infection, of the genitals as well as the organs most extensively affected. The frequently observed fact of the limitation within the internal os of tuberculous endometritis is forcibly noted. Another paper of interest is that of Dr. Edward Reynolds, of Boston, on "The Value of the Forceps in Complicated High Arrest of the Breech, with Report of Two Cases." The description of the adjustment of the blades to the breech is unusually instructive, inasmuch as the

safe application, in which the lower part of the breech is grasped, and that by the tips alone, will enable future operators to appreciate the proper application, whereas otherwise the instrument might be introduced too far, in the mistaken idea that a high application is the desid-

In addition to these papers that of the President, Dr. Byrne, of Brooklyn, upon the conservative versus the radical treatment of "Carcinoma of the Uterus," is im-

TABLETS FOR THE MEASUREMENT AND RECORDING OF THE MAGNIFIED IMAGE OF THE FUNDUS OF THE EYE. By Dr. FLAVEL B. TIFFANY, of Kansas City, Mo. Published by Osborne & Pitrat, Kansas City, Mo.

THE author of these ingenious tablets has taken advantage of the well-known fact that the enlargement of the upright image in ophthalmoscopy depends upon the distance to which the image is mentally projected. Assuming, therefore, that the magnification of twenty times-about the ordinary amount of enlargement in the upright image-necessitates a chart of such dimensions to be placed thirty centimeters back of the observed eye, the observer is at once given a graduated blank by which the details of any fundus oculi, in almost the exact size of those seen in the ophthalmoscopic picture, can be projected on a sheet by his unused

We have successfully taken advantage of the method in a number of cases, though we find by comparison with others that it does not adapt itself to those observers who are presbyopic and not ambidextrous.

THE DISEASES OF CHILDREN, MEDICAL AND SURGICAL. By HENRY ASHBY, M.D. Lond., F.R.C.P., and G. A. WRIGHT, B.A., M.B. Oxon., F.R.C.S. Eng. Second edition. Edited for American Students by WM. PERRY NORTHRUP, A.M., M.D. 8vo, pp. 773. New York and London: Longmans, Green & Co., 1893.

THE additions and comments of the American editor have not been included in the text, but are collated in the appendix with proper reference. The formulæ have been rewritten to conform with the U.S. P. The work itself has, by its passing to a second edition, been stamped with professional approval; and it is, indeed, an accurate and reliable guide to the diagnosis and treatment of the diseases studied. We note with pleasure the acknowledgment made by the authors to successive generations of house-surgeons whose well-kept notes of cases have been utilized. Few American authors can make similar acknowledgment. The style of the authors is easy, but clear; and while it is concise, no important fact has been sacrificed to condensation, We can cordially commend it to student and practitioner.

CORRESPONDENCE.

FRACTURE OF THE HUMERUS DURING DELIVERY.

To the Editor of THE MEDICAL NEWS,

SIR: I see reported in THE NEWS several cases of fracture of the femur during delivery. Allow me to report a case that occurred to me a good many years ago, and has never been reported.

I was engaged to attend Mrs. the wife of a mill-owner, a very stingy fellow. When labor commenced he sent for a "wise woman," who, he was told, knew more about such cases than any doctor, in order that he might save a part of (in those early days) my small fee. About daylight one morning he came for me at a hard gallop, saying that his wife was bleeding to death. I went at once, and found a rather severe case of post-partum hemorrhage, and a roomful of awfully frightened women, the "wise woman" as white as Hamlet's ghost, for her patient was in a dead faint. Happily the patient responded to stimulants, and I succeeded in arresting the hemorrhage.

The nurse then said to me: "Doctor, there is something wrong with the baby's arm." On examination I found a simple fracture of the humerus, between the upper and the middle third. I could not learn the presentation. I reduced the fracture, and put it up in a moulded pasteboard splint. Union took place in a short time, and there was no deformity.

I had the satisfaction of charging and collecting full fees in both cases. I also introduced the next baby. N. AGNEW, M.D.

Respectfully, WINNIPEG, MANITOBA, CANADA.

SOCIETY PROCEEDINGS.

AMERICAN MEDICAL ASSOCIATION. Forty-fourth Annual Meeting, held at Milwaukee, Wis.,

> June 6, 7, 8, and 9, 1893. (Continued from page 677.)

SECTION OF DISEASES OF CHILDREN.

FIRST DAY-JUNE 6TH.

DR. CHARLES G. JENNINGS, of Detroit, delivered the President's Address, which was entitled "Some Recent Advances in Pediatrics." After thanking the Section for the honor conferred upon him, the Chairman called attention to the recent changes in the organization of the Association that give to the Section work, and to the scientific work their proper position as leading factors in the organization. He pointed out that the Section has heretofore lacked cohesion and cooperation. The membership of an undifferentiated section is diffuse and unstable, and it cannot exert the influence in the Association and the profession that it should. Pediatrics covers almost the whole field of medical and surgical specialties. Including in its membership general practitioners and specialists, the Section is most favorably organized to carry out its function of collecting and disseminating the most advanced teachings in this department of medicine. The Chairman urged the necessity of more firmly establishing the Section's autonomy. He advised the cultivation of a more intimate social relation among the members, full and prompt reports of discussions and proceedings, and the arrangement by the officers of the section of short, practical papers, in series, on live subjects. The Chairman then briefly reviewed some of the recent advances in pediatrics. He called attention to the more extensive use of the bath in febrile diseases, and noted that the bath frequently, and properly, replaced active antipyretic drugs. Considerable attention was devoted to a review of the recent investigations into the etiology of diphtheria and the relation that the disease bears to the various pseudo-membranous throat-affections.

The difficulty was emphasized of accurately diagnosticating diphtheria from similar conditions by clinical means alone, and the necessity of bacteriologic examinations was urged. In cases of doubt, and when these examinations cannot be made, the doctrine of isolation should be observed to its farthest limits.

DR. W. A. DIXON, of Ripley, Ohio, read a paper entitled "Observations on Isolated Cases of Diphtheria." He stated diphtheria is a common and destructive disease, from which rich and poor alike suffer severely. Sanitary conditions seem to have little to do with its prevalence, It has always displayed a marked tendency to prevail in sparsely populated districts rather than in centers of population. In the United States it occurs as often on the hilltop as in the crowded city tenement houses. Pathologic reports are replete with statistics showing the communicability of diphtheria from the lower animals to man.

Cases of isolated diphtheria were cited to show that children have been infected from birds, fowls, pigeons, and cats. In 1865 an epidemic of malignant and fatal diphtheria prevailed in Southern Ohio following a period of great fatality among the hogs and chickens of that region.

In another case, a child, whose home was perfectly isolated, and who had a pet cat which it nursed continually, and which had been ill some days and had a discharge from the nostrils, was seized with diphtheria.

DR. G. BENSON DUNMIRE, of Philadelphia, read a paper entitled "Some Observations in Treating Cases of Diphtheria."

DR. F. E. WAXHAM, of Chicago, presented the following table of statistics of intubation:

	Age.		1	Cases.	1	Recoveries,	Percentage.
Under 1 y	rear			13		4	30.76
1	16			62		13	20.96
2 y	ears .			81		25	30.86
3	14		٠	85		32	37.64
4	16			90		35	38.88
5	61			43		19	44.18
6	41			26		7	26.92
7	46			29		10	33.33
8	44			13		8	61.53
9	54			7		3	42.85
10	64			7		3	42.85
11	64			1		I	100.00
12	**			3		0	00 00
13	16 .			1		0	00.00
. 14	61			1		0	00.00
20	44			1		0	00.00
36	44			x		0	00.00
43	44			1		1	100.00
60	41 .			1		0	00.00
				466		161	34.54

DR. WAXHAM also read a paper entitled "The Therapeutics of Diphtheria." He expressed the hope that the inoculation-treatment of diphtheria will soon replace all other methods of treatment. There are still great difficulties to overcome, however, before the method can be applied to man. The treatment of diphtheria may be considered under the following four heads: (1) Nourishment, (2) Stimulation, (3) Internal Medication, (4) Local Antisepsis. The importance of feeding is often overlooked, especially as to the amount that is taken. On account of loss of appetite, the patient may refuse all kinds of food. In such a case, peptonized food by enema or by the stomach-tube may be resorted to. Mild cases do not demand the use of alcoholic stimulants, but severe cases do demand them. When required, alcohol should be given freely at least one or two teaspoonfuls every hour, or half-hour, according to the urgency of the case. If necessary, strychnine, musk, or ammonia should be given. The tincture of ferric chlorid is a valuable agent. It should be given in frequent and full doses-ten, fifteen, or twenty drops to young children and should be repeated every hour or half-hour. Mercuric chlorid or potassium chlorate may be given, but in severe cases of diphtheria either is too irritating to the kidneys. The local antiseptic treatment is of great importance. Carbolic acid, mercuric chlorid, tannic acid, chlorin water, or hydrogen dioxid may be used. The results, however, will not depend so much upon the remedy employed as upon the method of employment. In the strength of 1 to 4000 mercuric chlorid is not irritating. Hydrogen dioxid, I to 4, is not irritating. The spray is to be preferred to the douche. One of the most efficient means of flushing the nasal cavities is by the use of the soft catheter. The indications in the treatment of diphtheria are to destroy the bacilli and to support the system by abundant nourishment, free stimulation, and full and frequent doses of iron.

DR. J. A. LARRABEE, of Louisville, stated that it is always well to give the patient the benefit of any doubt and treat the case from the first as though it were one of diphtheria. There are some points by which the diagnosis can often be made rather early. The constitutional disturbances of tonsillitis are sometimes profound, but they are not associated with the blood-changes observed in diphtheria. Albuminuria is present in almost all cases of diphtheria in the second stage. Glandular enlargements are also present in diphtheria. The appearance of the pseudo-membrane cannot be relied upon. In the preventive treatment tincture of ferric chlorid is to be preferred, which should also be given during the whole course of the disease. It is usually not given in large enough doses. It increases the number of red blood-corpuscles, and thus renders it possible for a larger amount of oxygen to be taken into the body. It also stimulates the appetite and acts as a diuretic. Local treatment should precede or at least accompany the constitutional treatment. Mercuric chlorid is objectionable. If used as a gargle or spray death may result. Hydrogen dioxid, used carefully, does well. Too strong solutions may cause sloughing. As it is almost impossible to spray or irrigate the child's throat the insufflation of boric acid and pepsin or papain is to be preferred. Argentic nitrate is also entitled to consideration. As one attack of diphtheria does not confer immunity to subsequent attack, the value of treatment by inoculation is doubtful.

DR. DOUGLASS, of Detroit, stated that if the tonsils are much inflamed and swollen and encroach upon the air-passages, it is well to incise them. He has not observed this operation to be followed by septic poisoning.

SECOND DAY-JUNE 7TH.

DR. W. S. CHRISTOPHER, of Chicago, read a paper on "The Pathogenesis of Bronchitis." It was shown that bronchitis arises from a large variety of causes. This multiplicity of cause indicates that the disease is probably always secondary and never primary. The principal causes are the poisons of acute infectious diseases, intestinal infection, malnutrition, and bronchoadenitis.

DR. C. L. DODGE, of Kingston, N. Y., read a paper on "The Pathology and Symptomatology of Acute Bronchitis and Broncho-pneumonia." The pathology of acute bronchitis was stated to be "an inflammation affecting a mucous structure, leading to a secretion of mucus and the production of muco-pus in greater or less abundance." The symptoms differ much in degree and as the disease is mild or severe. In addition to the usual premonitory symptoms of coryza, etc., there is a harsh paroxysmal cough. The respiration is accelerated, and in mild cases there is slight fever. In severe cases the fever is considerable and the pulse frequent. The temperature ranges from 101° to 102° F. In severe cases it may reach 103° F. The skin is dry and hot and the cheeks flushed. The cough in many cases is harsh, dry, and persistent. There is also more or less laryngitis, together with pharyngitis of mild grade.

Catarrhal pneumonia is not, like croupous pneumonia, a distinct and independent disease clinically, but in the great majority of cases it is a secondary phenomenon in the course of acute and chronic diseases of various kinds. It almost always follows bronchitis. The same process that causes catarrh of the bronchial mucous membrane, in its further course, invades the bronchioles and alveoli and leads to catarrhal pneumonia.

Broncho-pneumonia is usually preceded by acute bronchitis. The extension of the disease to the lungtissue proper manifests itself by dyspnea, a change in the character of the cough, and the purulent character of the sputum. When extensive collapse takes place the dyspnea increases, the temperature falls, the cough ceases, and the child rapidly sinks into a comatose condition. The pulse is extremely rapid and runs from 150 to 200. The respiration is also increased, and in severe cases reaches 80 per minute. The temperature will average 104° or 105° F. in acute cases. There is no regular ratio between pulse, temperature, and respiration. Anorexia and thirst are prominent symptoms. On auscultation during the early stages râles of all sorts and sizes may be heard, but it is at a later period that there are persistent subcrepitant râles in one or more

DR. J. M. G. CARTER, of Waukegan, Ill., read a paper on "Some Phases of Broncho-pneumonia in Children." Attention was directed to a class of cases that do not receive the careful observation that they merit. Common colds in children frequently develop into severe attacks of broncho-pneumonia. Many cases recover spontaneously, so that the number of cases that come under the observation of physicians is diminished, and the rate of mortality ascribed to broncho-pneumonia appears high. A recurrence of broncho-pneumonia from colds in adelescence and middle life may develop fibroid pneumonitis. In old people and in children under five years of

age, broncho-pneumonia causes the greatest mortality. In simple cases, the chest should be enveloped in cotton batting, the patient kept quiet in bed, and mild anodyne expectorants administered.

DR. F. S. CHURCHILL, of Chicago, read a paper on "The Pathology and Symptomatology of Croupous Pneumonia." Croupous pneumonia in the child, as in the adult, was defined as an acute, infectious, self-limited disease, having its chief pathologic manifestations in the lungs. The pneumonic lung of the child is not as greatly distended nor as solid as that of the adult. It is denser and darker-colored than normal. The surface is rather smooth, and the inflammatory exudation less marked than in the adult. It is not uncommon to find abscesses near the surface following the third stage.

The microscopic appearances are much the same as in the adult. The exudate is usually fibrinous, and contains numerous red and white blood-corpuscles. When it begins to liquefy it contains diplococci, which closely resemble the pneumonia-microbes. The morbid process has a strong tendency to affect the lower lobe; more frequently on the right than on the left side. The onset is usually sudden, with stabbing pain in the side, suppressed cough and rusty sputum; but in children this complex of symptoms is not always present, and the diagnosis is often difficult. The child usually complains of pain in the epigastrium, and headache, accompanied by rise of temperature. The child is usually found breathing rapidly with an "expiratory grunt," its face flushed, the alæ nasi dilated, and the expression painful. The affected side may be noticed to lag. Vocal fremitus may be increased. The respiratory murmur may be bronchial. Râles are rarely heard in children less than three years old. On percussion, a sense of resistance is appreciated over the affected area. Cerebral symptoms vary in intensity, from restlessness to severe convulsions. A certain degree of cyanosis may be present. The temperature varies from 102° to 106°. The pulse and respiration are frequent. The proportion of cases ending by crisis is greater than in adults.

DR. I. N. LOVE, of St. Louis, read a paper entitled "The Therapeutics of Bronchitis." He stated that some observers maintain that bronchitis is due to a special germ, whose port of entry is the air-passages, but the majority admit, however, that bronchitis is, as a rule, simply a result of "taking cold." The predisposing conditions are constipation, improper diet, disturbed digestion, perverted secretions, and exposure to cold. In the therapeutics of bronchitis, the general torpor of the glandular apparatus must not be ignored. The bowels should be thoroughly moved by means of calomel or a saline, followed, if necessary, by castor oil. These remedies mentioned, together with turpentine, serve to stimulate the secretory activity of the mucous glands of the air-passages, and favor expectoration. Local applications may be made to the chest. If the temperature is high, some one of the antipyretics should be given in a stimulating menstruum to reduce the temperature to proper limits. Too high a temperature is dangerous, no matter what the affection may be. It is to be borne in mind that bronchitis in children may terminate in catarrhal pneumonia. Stimulation at the proper time is important, and it is often of advantage to give one or two teaspoonfuls of equal parts of whiskey, glycerin, honey, and some preparation of malt. The secretions should be stimulated and rest secured. Dover's powder may be used to advantage. Malaria should never be overlooked, for it is remarkable how often it is a potent factor in a case of bronchitis in a child. After the acute stage has passed, turpentine, or terpin hydrate can be given, together with mild tonics and a nutritious diet. Special emphasis is to be laid upon elimination, tranquillization, stimulation, nutrition, and oxygenation.

DR. F. S. PARSONS, of Northampton, Mass., read a paper on "The Therapeutics of Broncho-pneumonia." He stated that a comfortable, well-ventilated room is essential to successful treatment. Exposure is to be prevented, if possible. The patient should be clad in soft flannel and kept in bed. It is well to direct a chest-protector to be worn. This is preferable to poultices, ointments, and other applications. If the case is seen early, medication is to be directed to the cough and to the promotion of resolution, but if seen late and the child is weak, stimulation must be resorted to, for which brandy is to be preferred. Digitalis in drop-doses is valuable. If the temperature is high acetanilid or aconite may be used; the latter being preferable in very young children. It may be necessary to administer an emetic to free the stomach and bronchial tubes. Under such conditions small doses of calomel may be administered to empty the stomach and intestines. The position of the child's body should be changed frequently, to prevent hypostatic congestion. If there is a tightness of the chest, the boiling of water in the room will frequently do good. To stimulate the action of the mucous glands and to increase the watery elements of the secretion the spray is indicated, and the best drug is ipecac in some form. Opium may be combined with ipecac to relieve irritability. As a diuretic, potassium citrate is to be recommended. Ammonium and other stimulating expectorants are useful in the later stages. During convalescence tonics of iron and a good nutritious diet are always

DR. J. A. LARABEE, of Louisville, read a paper on "The Therapeutics of Croupous Pneumonia." He recommended a formula containing spirit of nitrous ether, f 3 jss; potassium acetate, 3 jss; solution of ammonium acetate, water, and camphor, each, f 3iij. The solution should be slightly acid in reaction. The dose for an adult is one tablespoonful; for a child a teaspoonful, every two hours. To this may be added aconite or veratrum viride, and during convalescence tincture of ferric

Death usually occurs through the heart. Pari passu, with the consolidation of the lung, and in direct proportion to its extent, the transfer of blood from the right to the left heart is mechanically obstructed. The condition would be rendered still more dangerous by the use of such a stimulant as digitalis. The ligation of the lower extremities, sufficiently to prevent return of venous blood, relieves the heart. The application of dry cups and flaxseed poultices to the chest is of service. The best results are to be obtained, without jeopardizing the life of the patient, and without interfering with convalescence: (1) By agents that determine the blood to the skin; (2) by belladonna; (3) by nitro-glycerin and the nitrites; (4) by applications to the chest.

The mixtures usually prescribed for the cough of

pneumonia contain ipecac and squills to loosen, senega and ammonium to stimulate, wild cherry and other tannates to check secretion; alkalies to lessen the viscidity of the mucus; and opium to dull sensibility. Water given in abundance and at short intervals is the best and only expectorant required. Purgatives add to the general depression. If the bowels need attention, they should be moved by simple enemata.

Hyperpyrexia usually yields to the warm bath or pack.

The pack is preferable to the bath.

DR. JAMES B. HERRICK, of Chicago, stated that in cases of central pneumonia the physical signs are frequently so imperfectly marked that the diagnosis is very uncertain. The pain is so frequently referred to the epigastrium that attention is attracted to this region, but the rapidity of the respiration should direct attention to the chest. The broncho-pneumonia that occurs in the new-born is probably due to the inhalation of liquor

DR. E. E. GRAHAM, of Philadelphia, stated that if, in the treatment of pyrexia, he were limited to a single agency, he would not hesitate to accept cold as the best, If cold is applied at all, it should be done fearlessly. Depressing effects do not follow the proper employment of cold. Good results may also be obtained from cold water injections; from a pint to a pint and a half can be injected at a time. Aconite should be used with the greatest caution.

DR. CHARLES G. JENNINGS, of Detroit, stated that the bath is a stimulant to both respiration and circulation, as well as a good antipyretic. Drugs may be used in the beginning of the disease; but during the latter part of its course the bath, beginning at 90° and gradually reduced to 80° or 75°, acts as a stimulant to all of the functions and will reduce the temperature to a safe point.

Dr. J. M. G. Carter, of Waukegan, Ill., stated that, according to statistics, there are 561 deaths per million due to bronchial affections in the city, to 165 per million in the country. Among the Indians there are only 10 deaths per thousand attributable to bronchitis, while among the whites there are about 28 per thousand. These figures go to show that exposure does not play a very important rolê in its causation.

THIRD DAY-JUNE 8TH.

Dr. J. A. Work, of Elkhart, Ind., read a paper on "The Importance of Early Effective Elimination in the Zymotic Diseases of Children." All children equally exposed to a certain contagion do not alike contract disease. Because some are in a healthy condition, the system is in a normal physiologic condition, the poison finds no nidus of effete matter or no lodgment to encourage it to remain.

A system thoroughly cleansed of all effete and residual matter is rendered less susceptible to, if not entirely exempt from, the poison of scarlatina. It is, therefore, a rational procedure, in treating little patients, to eliminate as early as possible through all channels and maintain the normal action of the eliminative organs throughout the entire course of the disease.

Early elimination is important in all zymotic diseases, and, if practised early, will in many cases prove abortive.

DR. MARION THRASHER, of San Francisco, read a paper on "Dentition and Some of its Diseases,"

pointed out that normal dentition is comparatively harmless, but in this age of civilized refinement abnormal dentition is the rule, and the danger is constantly increasing. A few decades ago, when it was more general than now the practice of mothers to nurse their children, the fatality was not nearly so great.

Dentition in artifically-fed children develops incidentally many diseases. The commonest are irritative fever, indigestion, convulsions, stomatitis, capillary

bronchitis, and pneumonia.

Purulent conjunctivitis may arise when the upper canine teeth are being cut, the inflammation being transmitted through the antrum of Highmore. Cough is often present when the upper teeth are being cut. Eczema impetiginosum may also occur at this period. During the evolution of the teeth the submaxillary glands are often swollen and the cervical glands may enlarge and suppurate, especially in scrofulous or rachitic children.

The hygienic and dietetic conditions should be made most propitious. The diarrhea that is often present is usually best met with minute doses of hydrargyrum cum creta, sodium bicarbonate, and tannin, for a day or two, followed by oil. Excessive temperature should be controlled. Cutaneous affections require no special attention during this period. Diseases of the respiratory organs should receive attention. Restlessness should be overcome by hot baths, aconite and bromids.

Delayed dentition in puny children may be remedied by vigorous constitutional treatment, including pure air, proper food, bathing, ferric citrate, cod liver oil, quinine.

DR. A. FOSTER, of Chicago, laid stress upon the importance of discriminating between the effects of teething and those of improper feeding. He also pointed out that there may be a reflex relation between the cutting of teeth and the stomach, as well as there is between the uterus and the stomach, and that all of the trouble arising during dentition may not be due to the stomach alone.

DR. J. WELLINGTON BYERS, of Charlotte, N. C., read a paper entitled "Phenomena and Causes of Gastrointestinal Fever (not typhoid.") He said that there is a large class of gastro-intestinal disorders characterized by phenomena closely resembling those of typhoid fever, which deserve special attention and study from the fact that they are often confounded with the latter condition, and a line of treatment is instituted that serves only to aggravate and protract them. These disorders have been variously designated as gastric fever, gastro-intestinal fever, pseudo-typhoid, abortive typhoid fever, simple fever, and continued fever, though they are all phases or forms of the same disease, namely, disordered digestive processes leading to putrefactive changes in the alimentary canal, and the absoption of toxins into the circulation. They set in with a diversity of symptoms, in accordance with the individual attacked and the specific etiologic agents. They are most frequently traceable to errors of diet; but fatigue, anxiety, shock, undue physical and mental exertion, exposure to a foul, damp atmosphere, and the heat of the sun may act as causes. As a rule, young children and those with feeble digestive powers are most susceptible. The attack sets in with a feeling of indisposition, followed by chilliness, fever, headache, thirst, anorexia, and pain in the back, abdomen, and limbs. The tongue is coated with a

white fur, and the pulse is accelerated. There is commonly constipation, though in young children diarrhea is often present. The pyrexia reaches its maximum early on the first or second day. The duration of the attack is usually less than a week if proper treatment be instituted; otherwise, the disease may continue for an indefinite time, and thus simulate typhoid fever. As contrasted with the latter, the rise of temperature is sudden, the highest, 104°, being reached in the first few days of the invasion. The bowels are most frequently constipated; there is absence of the characteristic typhoid tongue; and there is no circumscribed iliac tenderness or tympanites, but a general soreness over the entire abdominal region. The prostration of the nervous system is not so marked as in typhoid, though patients often complain of great muscular weakness even after a few days' illness. In an etiologic connection recent developments in physiologic chemistry and bacteriology have shown that whenever from any cause the normal digestive functions are perverted, putrefactive processes occur in the alimentary canal. Disordered innervation, originating under debilitating circumstances of environment, acts as a predisposing cause by leading to perverted glandular and cellular function, thereby facilitating the production and absorption of alkaloidal poisons from the digestive tract. The old notion that a torpid liver is conducive to putrefactive changes in the intestines is the correct one. The hepatic condition is frequently the first recognizable stage in the causation of the disorders enumerated, and the first step in all treatment for their alleviation should be directed toward the restoration and maintenance of the hepatic functions. The first element of success is the differentiation from typhoid conditions. This being done, food should be withheld until the putrefactive substances have all been removed from the canal, and then a line of medication should be instituted that will tend to restore normal cell-function by promoting the trophic forces and the anabolic processes.

DR. J. SCHENCK, of Mt. Carmel, Ill., read a paper entitled "Cholera Infantum; Its Treatment in Malarial Localities." He defined cholera infantum as a disease of early childhood, in which there is exalted excitability of the nerve-centers and of their meninges, producing high fever and causing sudden attacks of forced vomiting and violent purging, which, if not controlled, result in exhaustion and death in a short space of time. The primary cause is continuous, high summer heat, both night and day, combined with malarial or other miasmatic poisoning. The secondary causes are indigestion and fermentation of the food-contents of the alimentary canal, including decomposition of the normal secretions and excretions of the digestive tract.

The preventive treatment is mental and physical quiet. The body should be kept as cool as possible during the time of greatest heat; pure and easily assimilable foods

should be given; all unhygienic conditions should be removed; digestion should be aided by pepsin; malarial influences counteracted by quinine, and nervous ex-

citement by physostigma venenosum.

For curative purposes, the little patient should be placed on a rubber bag filled with cold water; exalted nervous action should be inhibited with sodium bromid and physostigma, repeated until their physiologic effects are produced, or, in older children, hypodermatic injections of morphin may be given to accomplish the same object; cold egg-water should be given to allay thirst and fever; small doses of morphine and carbolic acid to control vomiting, and large mustard plasters should be applied to the extremities to relieve central congestion; the secretions and excretions should be stimulated by means of small doses of calomel and sodium bicarbonate. After reaction has taken place, the case is to be treated as one of severe entero-colitis—a condition that is almost certain to follow.

DR. M. G. SLOAN, of Dexter, Ia., read a paper entitled "Meningocele, with Case." The occurrence of pure meningocele is doubted by many, and is certainly very rare. The diagnosis is difficult. The condition is often diagnosticated as hydro-cephalocele. With modern antiseptic methods the operation for removal should be a comparatively safe one. In the case reported and operated upon the result was a perfect cure. Not a vestige of anything resembling brain-tissue was found.

DR. THOS. H. MANLEY, New York City, read a paper entitled "The Etiology, Morbid Anatomy, Diagnosis, and Treatment of Infantile Hernia of the Inguinal Type in the Male." It was maintained that the predominating elements in infantile hernia are anatomic, but heredity also plays an important rôle. The mal-descent of the testis is more often responsible than all the other known causes combined. The causes being so variable, the treatment also must be variable. The diagnosis of the inguinal type of infantile hernia is not difficult in the majority of cases, but in many forms of cystic disease of the cord an accurate diagnosis is not always possible.

The treatment will include attention to diet, clothing, rest, support or pressure, removal of such causes as occasion straining, and radical operation.

DR. WILLIAM E. WIRT, of Cleveland, Ohio, read a paper entitled "Hernia in Children." He spoke of the frequency and varieties of hernia in children. The condition is much more common in males than in females. Umbilical and femoral hernia are much more frequent in females than in males, but femoral hernia is rare in children. As etiologic factors, climate and the condition of the patient were mentioned. The treatment was considered under three heads, viz.: I. General treatment; 2. Mechanical support; 3. Operative measures.

A paper entitled "Acute Endocarditis of Children; Etiology, Symptomatology, and Treatment," by C. N. HIGHLEY, of Conshohocken, Pa., and one entitled "Primary Syphilis and Gonorrhea in Children," by B. M. RICKETTS, Cincinnati, Ohio, were read by title.

(To be continued.)

AMERICAN ACADEMY OF MEDICINE.

Eighteenth Annual Meeting, held at Milwaukee, Wis., June 3 and 5, 1893.

(Continued from page 680.)

SECOND DAY-JUNE 5TH.

DR. BENJAMIN LEE, of Philadelphia, presented a paper entitled "The Duty of the State to Medicine." He urged that the State should recognize that it incurs obligations at the same time that it imposes obligations upon the medical profession. Among these are the

duty of according to physicians occupying official positions equal recognition, rights, privileges, ranks, and emoluments with others, notably the legal and military professions; of securing to the medical profession such a standard of education and attainments as shall enable it to command the respect due it; of paying for services demanded; of protecting its members against malicious prosecutions; of officially recognizing achievements in medicine and surgery, and, finally, of recognizing the importance of State medicine. "The health of the people is the first duty of the statesman."

DR, GASTON said that the public should be compelled to recognize its obligations to the profession by statutory enactments, if necessary. The public seems more ready to recognize the quack than the educated physician. Medical officers should be empowered to perform their duties without secondary orders from superior officers. The necessity of preliminary training was emphasized. Dr. Gaston deplored the tendency in the South to neglect such training. Without preliminary education the student cannot understand medical lectures, and part of this training should be under preceptors and in active office-work. The necessity of the business aspect of the physician's life was urged as an element of success. The State should be required to pay for all medical services demanded. Pay for expert testimony by the courts is at present a farce. The profession should have a cabinet officer. If the profession would work together. this could be brought about. The organization of State Medical Examining Boards is a wise choice of evils. School diplomas cannot be relied upon. In malpractice suits the innocent should not be allowed to suffer, Rashness and ignorance should be punished.

DR. MARCY called attention to the fact that by the proposed extension the work of the Academy would be enlarged. The physician is not only a doctor; he is also a man and a citizen. The coroner system, for example, is a blot and a disgrace to our civilization.

DR. CONNOR said that if every physician taught his patients and friends their duties, there would be no trouble. The teachers do not stand together. Most malpractice suits originate with some member of the profession. The profession should be the teacher.

DR. Holmes alluded to the injustice of hospitals not paying physicians, as every other attendant—nurse, superintendent, etc.—is paid. The remuneration of the physician may consist in permission to utilize clinical material for clinical instruction. The reimbursement would come back to the public. It is an axiom, or should be, that no gratuitous service can be demanded of a physician. If he does give it, it is given as a delegate from the profession.

DR. C. C. BOMBAUGH, of Baltimore, read an excellent paper entitled "Literary Recreations of the Physician." Among other things, he said that there must be some form of recreation to compensate for the trials and anxieties of the physician, and among the methods of relaxation one is in weaving from his own unfettered fancies story or song, poem or play, essay or critique, brochure or quarto. The Index-Catalogue of the Surgeon-General's Office shows what medical men have done for medical literature. While they have not been so prolific in general literature, they have contributed much that commands widespread admiration. A great deal of this, it

is true, has been written by those who have withdrawn from active practice, hence Copernicus, Galileo, Servetus, and others of their class are not thought of at the first as physicians.

Others have found themselves not able to bear the wear and tear of active practice. Among these may be found Schiller, Thomas Huxley, William Farr, Erasmus Darwin, and Charles Lever. Others, like Abraham Cowley, never attempted to practise, but still others mingled their literary pursuits with an active practice, Mark Akenside, Dr. Arbuthnot in the reign of Queen Anne, Sir Richard Mead, who was physician in ordinary to George II, were mentioned among others.

America has quite a respectable list of such men. from Benjamin Rush, David Ramsay, Samuel Hopkins, and others of the Revolutionary period, to authors of the present day. Literary occupation is not work, however, in the ordinary sense of work; it is diversion, the safetyvalve for weariness. Thus the Autocrat of the Breakfast Table has shown how the Anatomist in the Lectureroom could trace the tortuous windings of the trigeminus, and then go home and pen such immortal verse as the "Living Temple" or "The Chambered Nautilus." Fair characterization has not yet been accorded the physician by the modern novelist. The satire of the earlier writers has castigated the pretenders of the past, but the qualities and qualifications of the end-ofthe-century physician, whose education ends only with his life, who meets conscientiously all moral and legal responsibilities, or who is ever ready for self-renunciation and self-sacrifice, have not been fairly weighed and measured by the end-of-the-century literary class.

Rudolph Virchow, who has not only reached the top of the ladder as a biologist and a pathologist, but has also climbed high as a statesman and a politico-economist, recently remarked that, "after wearying, useless efforts in political, social, or religious matters, scientific work is a recreation."

DR. J. CHESTON MORRIS, of Philadelphia, presented a communication entitled "The Care of Abandoned Infants." He said that in Roman times the infant born in the house was brought to the head of the house, to be acknowledged and taken care of or to be refused and slain.

In medieval times, children abandoned by their parents were exposed in front of the church, for adoption. Gradually the foundling asylum took the place of this cruder method, until the horrible results in the way of infant-mortality and of incentive to criminal life caused the system to be abolished in France and Belgium.

In the United States the almshouse afforded a refuge not much better than the foundling asylum abroad, and few infants survived the troubles and diseases of infance.

In consequence of the publicity given to the facts some ten or twelve years ago, an institution called "Sheltering Arms" was established in Philadelphia, on the principle of trying to save the mother through the love for her offspring. The results have been most satisfactory and encouraging. No infant is refused, but the effort is always made to find and keep together mother and child. After several months of residence in the house, where the mothers do the most of the work.

a place is found in the country for her, and the effort is made to keep in touch with her by correspondence.

No woman is destitute wholly of the true maternal instinct, It may be clouded or overshadowed, and if she is permitted to make away with her child, or even to abandon it, her moral nature receives a wound difficult to heal and always leaving a horrible scar.

DR. C. McIntire, of Easton, Pa., read a paper entitled "The Importance of the Study of Medical Sociology." He quoted from the Century dictionary the definition of "sociology" as: "The science of social phenomena; the science which investigates the laws regulating human society; the science which treats of the general structure of society, the laws of its development, the progress of civilization, and all that relates to society."

Is there a special department of medical sociology? Are physicians peculiar, because physicians? If this view is taken, reference may be made to the past and present discussions on the codes of ethics. But there is the other view—the relation of the physician, as a physician, to the world about him. Medical sociology has, then, a twofold aspect—the social science of physicians regarded as a class, and the peculiar relations that the practice of medicine may bear to the social structure as a whole.

Does the physician differ from his fellow-men engaged in other pursuits? There is a language peculiar to the physician; there are customs peculiar to him, Other characteristics are a tendency to undue specialism and an incongruous admixture of conservatism and credulity. The relation of the physician to society at large is shown by his efforts to relieve suffering among the "under half"; in the results of his labor in sanitary science as a factor in the solution of municipal questions, the construction of dwellings; in the care of armies and of fleets; in the true presentation of the desirability of physical education. The classification suggested for sociology was that of Mr. Melville Dewey, which has been adopted in many of our large libraries. It includes statistics, political science, political economy, law, administration, associations and institutions, education, commerce and communication, and customs and costumes. Whenever medical knowledge is necessary to the elucidation of questions coming under either of these divisions, there are to be found problems of medical

The Academy has been devoting its energies to one class of educational problems, but this has so far advanced toward solution that it does not now require the entire effort of the Academy, and it will be well for it to enter this territory, at present unoccupied by any medical society.

The importance of the study of medical sociology may be seen from a consideration of the importance of some of the questions that ought to be discussed. The "political powerlessness of the medical profession" is acknowledged by all. As physicians usually endeavor to have legislation for the benefit of the community, and not for personal ends, this fact is to be deplored; and if the profession could wield the political power that it ought, much could be done for the common weal. The problem of the prevention of pauperism is one of the most pressing of the age and involves medical ques-

tions. Not the least among them is use and abuse of the free dispensary.

DR. Connor thought it wise to extend the same study to medical men as to bacteria, etc. Medical men have a great burden of secrets; they are, however, the silent men of the community. The medical profession thus differs greatly from other professions and is thus at a disadvantage. The relations of medical men to each other are peculiar and beyond the comprehension of the outside world. The medical man, indeed, is one of the pilots of life. In the lines of education these problems of medical sociology have a place. The study of the Constitution and the ideal arrangement would unite the profession into a band of brothers, working for the world, and based, so far as association goes, on scientific work.

In discussing the paper of Dr. Gould, DR. BAYARD HOLMES made a plea for a closer study of our requirements for endowment. How much will it require to run, in a first-class educational manner, a medical school with a four-year course and four hundred students, or forty students each with a good preliminary education? In one of the large cities it will cost fifteen thousand dollars a year, not including a dollar of salary to teachers. If in computing the salaries for instructors the standard of four recitation-hours daily in each of four years be taken, there would be sixteen recitation-hours that must be provided for. As about two-thirds of this work is either laboratory or clinical work, at least ten additional hours must be allowed, i.e., twenty-six hours of the teachers' time will be consumed. An average day's work for a college teacher is two hours. It would then require at least thirteen teachers devoting full time to teaching. The salary of these men at \$3000 a year would be \$30,000, which, if added to the running expenses, makes a total expense of \$54,000 a year for the ideal medical school conducted for education only. It may be said that clinical teachers should not be paid, but it is desirable that every teacher should be paid moderately.

There is now a great endowment going to waste or being perverted. The public pays for the care of the poor in public hospitals, and yet the cases are not used for educational purposes. Every free hospital should be a training-school for medical men and medical students. The public deserves this, and the patients would receive

The noble endowment that has been given hospitals in the name of charity is not always well directed. Thus, the extravagantly constructed Johns Hopkins Hospital costs more to keep in repair than would suffice to run a medical school. These great hospitals pauperize the people, pervert the legitimate rewards of the medical profession, render it so niggardly paid that the educated and refined cannot find in it a life-work.

The universities should be called on to do as well by medicine as they do by other schools. This is not

The following officers were elected for the coming year: President, Dr. George M. Gould, of Philadelphia; First Vice-President, Dr. C. C. Bombaugh, of Baltimore; Second Vice-President, Dr. J. McF. Gaston, of Atlanta, Ga.; Third Vice-President, Dr. S. W. French, of Milwaukee, Wis.; Fourth Vice-President, Dr. C. W. Foster, of Woodyard, Me.; Secretary and Treasurer, Dr. Charles McIntire, of Easton, Pa.

NEWS ITEMS.

Fatal Quackery .- A terrible story comes from a Hunarian city. A family of eight persons were suffering from scabies, and a wise woman was consulted, who sold them a strong mercurial ointment, with which the whole body was to be freely rubbed. The ointment, made of putrid fat, set up septicemia in a few days, and when a medical man was called in two children were already dead and a third lay dying. It was further stated that the remaining members of the family would also probably succumb.-Medical Press and Circular.

Pasteur Institute in the Punjab .- It is proposed to establish a Pasteur Institute in the Punjab Himalayas for the treatment of hydrophobia and for scientific research. To make available the benefits of the Institute to the whole of India, patients are, when necessary, to be conveyed free of charge from any part of India to the Institute.

Paul Guttmann, Sanitary Councillor and Director of the Hospital Moabit, died at Berlin on May 24th, of pneumonia complicating influenza, at the age of fifty-nine

James Moleschott, the distinguished physiologist, comparative anatomist, and anthropologist, died on May 20th, at Rome, of erysipelas, aged seventy years.

Mr. Marcus Beck, Professor of Surgery at University College, London, died May 21st, of diabetes.

BOOKS AND PAMPHLETS RECEIVED.

How to Operate for Hemorrhoids. By Charles B. Kelsey, M.D. Reprinted from the Therapeutic Gazette, 1893.

Free Incision of Abscess of Ostitis of Hip, and Closure without Drainage. By H. Augustus Wilson, M.D. Reprinted from the Transactions of the Philadelphia County Medical Society, 1893.

Practical Details in the Preparation of Plaster-of-Paris Bandges. By H. Augustus Wilson, M.D. Reprinted from the Philadelphia Polyclinic, 1893.

A Clinical Lecture on the Prevention of Idiopathic Rotary Lateral Curvature of the Spine. By H. Augustus Wilson, M.D. Reprinted from Annals of Gynecology and Pediatry, 1893.

The Influenza Bacillus. By J. A. Comstock, M.D. Reprinted from the Southern California Practitioner, 1893.

The Creation of God. By Jacob Hartmann, M.D. New York:

The Truth Seeker Company, 1893.

Metatarsalgia (Morton's Painful Affection of the Foot), with an Account of Six Cases Cured by Operation. By Thomas S. K. Morton, M.D. Reprinted from the Transactions of the Philadelphia Academy of Surgery, 1893.

The Internal Treatment of Lupus Erythematosus with Phosphorus. By L. Duncan Bulkley, A.M., M.D. Reprinted from the American Journal of the Medical Sciences, 1893.

Clinical Study and Analysis of 1000 Cases of Psoriasis. By L. Duncan Bulkley, A.M., M.D. Reprinted from the Maryland Medical Journal, 1891.

Third Biennial Report on Vital Statistics of the State of Minnesota for the Years 1890-91. Minneapolis: Harrison & Smith, Printers, 1893.

Certain Problems in Abdominal Surgery, Based on One Hundred Celiotomies done at the Kensington Hospital for Women. By Charles P. Noble, M.D. Reprinted from the Transactions of the Philadelphia County Medical Society, 1893.

INDEX.

ABBOTT, considerations concerning Asiatic | Aneurismthe etiology of membranous rhinitis, Abortion, curet in, 207 Abrams, pulmonary atelectasis as a cause of anemia, 293
Abscess, hepatic, suppurative pylephlebitis and, secondary to appendicitis, 397 ischio-rectal, 120 mastoid, 203 Abstractors, appeal to the pity of, 413 Abuses, professional, 302 Achillodynia, 187, 331 Actinomycosis of the lip, 216 prognosis of, 78 Advertisement, cheap, 195 Advertiser, sneak, 221 truthful, 248 Agnew, D. Hayes, Memorial Children's Ward, 248 N., fracture of the humerus during delivery, 697
Albumin in urine, delicate test for the presence of, 328
Albuminuria in the apparently healthy, 341
nephritis without, 391 Albumose, 208 Alcohol, tax on, 193 to allay the craving for, 692 treatment of diseases without, 553 Alcoholic liquors and tobacco, should they be furnished to prisoners? 443 Alcoholism, some personal observations and Alcoholism, some personal observations and reflections upon, 90

"Allopathic Czar parties," 388

Amber oil as a rubefacient, 466

Amblyopia of diabetics, 302
post-marital, 328

Amenities, professional, 389

"American Family Physician's Company," 194
Medical Association, address of the president of, 617
Milwankee, 664 meeting at Milwaukee, 664 proposed constitution and bylaws of, 388, 443 United, 443 organizations, the work of, 627 nervous, duration of life of, 229 Amputation of lower part of thigh, 548 Amusia, 525
Amyl nitrite, recovery after swallowing a teaspoonful of, 544 Analgen, 190
Ancrum, combined tonsillotome and uvulatome, 14 Anemia and chlorosis, relative utility of the various preparations of iron in the treatment of, 356 essential, etiology of, 161 for constipation attended with, 272 pulmonary atelectasis as a cause of, 293 Anesthesia, mechanism of, 49 psychic, for touch, 101

Anesthetization, danger of, by operator, 524 Aneurism of a sinus of Valsalva, 200

of the aorta, associated with insuffi-ciency of the aortic valves, 297 thoracic, venesection in, 520 Angina pectoris, 615 epilepsy or, 27 pseudo-diphtheric, relation of, to diphtheria, 17 Angio-neurotic edema, 381
Annals of Ophthalmology and Otology, 498 Anosmia, psychic, 101 Anthrax, cerebral, 131 Anticholerin, treatment of cholera with, 48 Antipyrin, peculiar toxic symptoms from, Antipyrin-poisoning, 18 Antiseptic adhesive application, 48 Appendicitis, conservative surgery in, 615 early operation for, 27
occurring on the seventh day following
labor; rupture into the bowel, 515

some recent cases of, 225 suppurative pylephlebitis and hepatic abscess secondary to, 397 Appendix, vermiform, incarceration of the

small intestine by, 244
Arey, practical helps in the use of the metric

system, 362
Armour, the practice of medicine in Ontario, Canada, 528 Armstrong, should we treat fever? 141

Arrenic as a prophylactic, 162
for epithelioma, 244
Arsenical poisoning, with special reference
to its domestic sources, 589

Ascites of hepatic origin, potassium bitartrate for, 162 treatment of, by faradization of the abdomen, 205

Aseptizable, 248 Ashton, angio-neurotic edema; a report of three cases showing some interesting manifestations, 381

Asthma, for, 300 spasmodic, paraldehyde for, 187 Astigmatism, a temporary change in the axis of, 598 Ataxia, Friedreich's, the etiology of, 695

Atelectasis, pulmonary, as a cause of ane mia, 293 Attendants, trained, 413

Audi alteram partem, 135
Ayres, a case of puerperal eclampsia at the seventh month, with a few thoughts as to treatment from practical experience, 212

BACILLUS, additional, found in water, 244 aërogenes meningitidis, 695 Bacteria of the intestinal tract, 550 Bacteriology in India, 472 Baldness, premature, causes and treatment Ball, final examinations, 530

Ballard, an improved antiseptic wire-suture twister, 354

Balloch, on the possibility of malignant de-generation in benign growths, I Barber-shop, dangers of, 577 Barclay, closure of the ear by growths of

bone; removal of bony obstruction; cure of otorrhea and of deafness. Three cases; description of operations; instruments, etc., 423

Barham, neurotic eczema, 317 Barium chlorid for epilepsy, 48 Bartholow, Roberts, 82 observations on the treatment of ascites

by faradization of the abdomen, 205 the treatment of cholera, 113

Baruch, hydriatic or hydriatric, 28 Beri-beri, the microbe of, 275

Berlin sewage farms, 359
Beta-naphthol, impure, the identity of "hydro-naphthol" and, 348
Bevan, disposal of sewage, 8
Biliary coloring-matter in the urine, a deli-

cate test for the presence of, 328
Billings's, Dr., estimate of the medical pro-fession as a calling, 221
Biology, pathology and, 360

Bishop, a new measurement in the study of fever, 99

Black, removal of the drum-head and ossicles in diseases of the middle ear,

tongue, 275
Bladder, ergot for atony of, 162
hernia of, 356
tuberculosis of, 457

urinary, seven cases of rupture of, 197 Bladder-wall, vesical calculus inclosing a

Bladder-wall, vesical calculus inclosing a portion of, 438
Blind, a type-writer for, 360
Blindness, psychic, in its variations, 101
Blood, to prevent the entrance of, into the mouth, in operations upon the face, 494
Blood-corpuscles, white, 273
Books, medical, the making of, 599
Booth, Dr., 414
Bowel, obstruction of, from Meckel's diverticulum 45

ticulum, 45 Bradley, three unusual fractures, 379 Brain, cystic tumor of, successfully operated

upon, 160
Briggs, antiseptic cataphoresis in the treat-ment of diseases of the uterus and its

appendages, 65 Bromoform-intoxication, 692 Bronchial catarrh, terpene hydrate for, 384 Bronchitis, chronic, and emphysema, for,

410

topical treatment of, 267
Bronchocele, myxedema following, cured by thyroid therapy, 438
Brucine and strychnine, a comparative study of the physiologic actions of, 369
Bryant, fracture of patella treated by continuous extension; patients not confined

to bed, 5 Buchanan, chemistry and therapeutics of nitrous oxid, 393
preventive treatment of tetanus, 110

Bureau of Medicine and Surgery of the Navy, 470 Burgess, prenatal closure of the pulmonary artery, 437
Burnett, partial myringectomy and removal of the incus and stapes for the relief of the lesions of chronic catarrh and otitis media, Butterine in the State of New York, 248 CAFFERY, secondary celiotomy following an oöphorectomy, 542 Cahill, summer diarrhea in children, 658 Calcaneus, talipes, talipes valgus and, 269
Calcium chlorid for pneumonia, 466
Calculus, vesical, enclosing a portion of the bladder-wall, 438 Calvarium, perforating epithelioma of, in a girl of fourteen, successful removal of, 242 Camphoid, a substitute for collodion, 410
"Canadian Practitioner," 248
"original communications" in, 414 Cancer-cure, 167 wood-sorrel as a, 363
Cancrum oris, with recovery, 493
Cantrell, irritation from local use of iodoform, 13 Capp, the treatment of carbuncle, 159 Caput coli, inflammation about, 561 Carbolic acid as an antiseptic, 220 poisoning from the use of, employed for expulsion of oxyures vermiculares, 158
Carbuncle, abortive free crucial incision in the treatment of, 307 treatment of, 159
Carcinoma, increase of mortality from, 304
in females, relative localization of, 306 parasitism of, 47 sporozoa and their relation to, 19 ood-sorrel for, 363 Cartledge, the present status of drainage in surgery, 124 Cataphoresis, antiseptic, in the treatment of diseases of the uterus and its appendages, Celiotomy, secondary, following an oophorectomy, 542 Central Hospital of Philadelphia, 332 Cesarean hysterectomy in a malacosteon, 354 Chambers, early operation for appendicitis, Chancroid, for, 132, 216 Chapin, the Gross Monument Fund, 390 "Charlotte Medical Journal," 381 Chicago, four years' course in, 577
sanitary condition of, 386
Lancet's investigation of, 413 Children, aural vertigo in, 409 Chloralose, 216, 357 Chloroform, a new study of the action of, 329 and the heart, 53 the use of, as an anesthetic, 374 Chlorosis, for, 520 the relative utility of the various prep-arations of iron in the treatment of anemia and, 356 Cholecystitis simulating intestinal obstruction, 383

Cholecystotomy, 45

stones, 296

Congress, 471 bulletins, 471

concerning, 361 excreta, disposal of typhoid and, 359 germs, experiments with, 333

Curvature of the spine, lateral, new aid in Cholerain Europe, 443 in Minnesota, prevention of, 333 news, dearth of, 360 our preparations for, 385 the diagnosis of, 452 Cutaneous disease, persistent stigmata re-sulting from, 281 protective inoculation against, 525 quinine for, 520 regulations at foreign ports to prevent, therapeutics of, 522 transmission of, through the interme-diation of flies, 328 treatment of, 113 with anticholerin, 48 winter, in Russia, 414 Cholesteatoma of the ear, 253 Chorea, etiology of, 429 for, 662 quinine for, 466 Choreoid movements in adult male, probably of hysterical origin; relationship to habit chorea, 231 Christian science dupers and dupes, 471 Cleanliness, municipal, 332 Club-foot, treatment of severe, 383 Cocaine phenate as a local anesthestic, 345 Code, no, 220 of ethics, revision of, 105 report of the committee on the revision of, 693
Cod-liver oil, emulsion of, 356
Norse mythology and, 276
Cohen, a case of uremia in an alcoholic patient with visceral complications successfully treated by venesection, diaphoresis, and nitro-glycerin, 298 notes on acute tonsillitis, 461 therapeutic notes, 351, 596 two cases of membranous enteritis, 156 College, medical, consolidation, 194 Colleges, medical, the consolidation of, 303 Colpoporrhexis, severance of the cervix from the body of the uterus, 243 Comedones, for, 300
Comma-bacilli, vitality of, 328
Complexion specialists in Berlin, 577
Confidence, professional, 695
Confidence, sanctity of, reposed by patients in physicians, 439 Constipation attended with anemia, for, 272 painful, of plumbism, for the, 574 treatment of, by means of rectal injections of oil, 495 Consultation with gentlemen, 249 Cooke, the use of chloroform as an anesthetic, with a case, 374 Copeland, venesection in rigidity of the os uteri and in allied conditions that may retard labor, 305 Cord, spinal, traumatic lesions of, 288 Coroners, scandal-mongering, 333 Cough of nasal origin, with report of a case, Cramp, writer's, treatment of, 410 Craniectomy and age, 476, 531 Cranium, denuded, operation for re-covering, 239 Creasote, iodoform, and guaiacol, to de-odorize, 300 treatment of pulmonary tuberculosis, 132 Credit to whom credit is due, 525 Cremation at Tokio, 611 with recovery after removal of gall-Virchow on, 400 Cholera, 440
and the water-supply, 332
Asiatic, considerations concerning, 622 Cretinism, sporadic, successfully treated by thyroid feeding, 662 Crime in England, decrease of, 414 Croton-water contamination, New York Croton-water contamination, New York
Academy of Medicine and, 303
Croup, diphtheric, case of, in which a
tracheotomy-tube was worn for sixty Asiatica in Boston, precautions against, at the Eleventh International Medical days, 691 Cure-alls in the daily papers, 359 Curet, Gottstein's, a modification of, 382 in abortion, 207

Cystitis, for purulent, 187 Cytotaxis, 575 DABNEY, clinical reports in laryngology, 492 Da Costa, relapses of typhoid fever, 154 Davidson, fracture of the femur during delivery, 614 Davis, treatment of puerperal eclampsia, 518 Death, stating the cause of, 359 under ether of Col. Shepard, 358 Death-rate in New York, 443 low English, 359
Deglutition, action of epiglottis during, 549
Delirium, for acute maniacal, 487 tremens, treatment of, 356 Dentists in England, 443
Dermatitis medicamentosa from iodoform, potassium iodid, and aristol, 240 Dermoid cyst, unusual finding in, 666 Diabetes mellitus following incised wound of head, 28 jambul in treatment of, 46 use of soya beans in, 662 of pancreatic origin, 275
pancreatic juice for, 216
Diabetic coma. 134
Diabetics, amblyopia of, 302
Diarrhea, summer, in children, 658
Diazo-reaction, additional notes on, 569 Diazo-reaction, additional to the blood of the blood of ferric chlorid in the treatment of, 162 ferric chlorid in the treatment of, 162 ferric chlorid in the treatment of, 162 ferric chlorid in the blood of the bl protective substances in the blood of convalescents from, 438 ravages of, 137 the relation of pseudo-diphtheric angina to, 17
without membrane, 552
Diphtheric croup, case of, in which a tracheotomy-tube was worn for sixty days, 6qI Dipsomania, bichlorid of gold treatment of, Disease, parasitism of, 165 Disinfectants, some commercial, analytic notes on, 595
Disinfection by high temperature, 410
Dislocation, subcoracoid, of the left shoulderjoint, 457
Dispensaries, New York City, statistics of the work of, 55 Dispensary abuse, 360 abuses, a lay view of, 220 conducted by women, 414 Diuretic and heart-tonic, 384 Dock, gonorrhea of the rectum, 325 Doctor, G. T. R., 359 Drainage in surgery, present status of, 124 Dropper and hypodermatic syringe, com-bination of, with improvements on both, Drum-head, removal of, and ossicles, in diseases of the middle ear, 400 Dulaney, traumatic empyema cating with the bowel, 689 Dulles, a splint for fractures and luxations at the elbow-joint, 214 Dupe, a willing, 525 Dura mater, sarcoma of, 519 Dwellings, model, for the poor, 136 Dyspepsia of hyperacidity, 565 Dyspnea, the nystagmus of, 627 Dysuria of senility, 412 EAR, cholesteatoma of, 253 closure of, by growths of bone, 423 influence of diseases of, upon the mental and physical development of the

child, 372

middle, removal of the drum-head and ossicles in diseases of, 400 Fareckson, purpura hemorrhagica, 10 Echinococcus cyst, large abdominal, 365 Echo from the past, 414
Eclampsia gravidarum et parturientium, in children induced by hernia of the cecum in the inguinal canal, 626 puerperal at the seventh month, 212 puerperal, treatment of, 518 Economy, expensive, 137 Ecphyaditis, 695 Eczema, neurotic, 317 Edema, angio-neurotic, 381 Education, higher medical, condition of, 136 endowed colleges and, 137 medical, 195 brains and money in, 553 of missionaries, 195
of women, history of, 413
of the deaf and dumb in England, 333
Eichberg, some experiments with modified tuberculin, 85 Elbow-joint, a splint for fractures and luxations at, 214 Electrolysis, epithelioma of the tongue cured by, 384 Embolism, pulmonary, from mercurial injections, 18 Empyema, traumatic, communicating with the bowel, 689 Endocarditis, chronic intermittent fever of, England's drink-bill, 304 Enteric fever without pyrexia Enteritis, membranous, for, 466 two cases of, 156 Epiglottis, action of, during deglutition, 549 Epilepsy, 489 barium chlorid for, 48 for, 565 or angina pectoris? 277 Epileptic colony in New York, 166 sleep-seizures, periodic, 598 Epithelioma, 433
arsenic for, 244
of the tongue cured by electrolysis, 384 perforating, of the calvarium in a girl of fourteen, successful removal of, 242 Ergot for atony of the bladder, 162
Erysipelas genitalium, due to the use of infected ointment, 148 Eshner, achillodynia, 187 Eskridge, cases of moral imbecility, of the opium-habit and of feigning, in which forgery is the offence committed, 29 traumatic myelitis, 235 Esophageal stricture and gastrostomy, 294
Esophagotomy, external, for the removal
of a foreign body from the esophagus, 189 in a child, 330 Esophagus, gastrotomy for the removal of a peach-stone from, 189 rupture of, 519 Esteem of one's fellows, 360 Ether, newspaper reports as to deaths from, 470
Etiquet, medical, 247
Evans, Keeleyism and Keeley methods, with some statistics, 477
periodic insanity in which the exciting cause appears to be the menstru function; report of a typical case, 538 Examinations, final, 530 Exanthems, medicamentous, 498
Excelsior College of Medicine, 221 Exophthalmos, pulsatile, 626

Exothyropexy, 406
Eye, indications for the enucleation of, 465

sclerosis, 355

peculiar injury of, 573 symptoms, early, of posterior spinal

sterni, 241

Groff, village and emergency hospitals, 533 Gross, Samuel D., monument fund, 276, 390

FACIAL nerve, peripheral palsy of, position | Gruwell, fracture of femur during delivery, of tongue in, 18
Faradic current, reduction of goiter by, 55 531 Guaiacol, iodoform, creasote, and, to de-Faradization of the abdomen, treatment odorize, 300 ascites by, 205 Fast, fifty days, 137 Gynecology among the insane, 276 Femur, fracture of, during delivery, 323, 531, HABIT-CHOREA, choreoid movements in an adult male, probably of hysterical origin, relationship to, 231 Hahnemannian tendencies and practices, 79 614 reason why neck of, does not break, 97 traumatic separation of lower epiphysis Fever, mountain, 376 Hamilton, Chas. S., report of operations new measurement in the study of, 99 should we treat, 141 upon hernias, 95 John B., gangrene of great toe; sub-coracold dislocation of left shoulder-Fisher, albumose, 208 Ford, legislation in reference to quarantine joint; tuberculosis of bladder, 457 Hand, apical pneumonia, 549 and sanitation at the port of Philadelphia, 473 sanitation in Philadelphia, 501 Hansell, prominent symptoms of hyper-phoria, as illustrated by thirteen successive cases, 181 Hart, medical journalism, 653 Forgery is the offence committed, cases of moral imbeelility, of the opium-habit, and of feigning, in which, 29
Foshay, some notes upon urinalysis, 201
Fracture, severe, of the vertex, with replacement of a large fragment of bone, 185 Hawkins, poisoning from the use of enemata of carbolic acid for expulsion of oxyures vermiculares, 158 Haynes, antiseptic midwifery, 28 Hays, cystic degeneration of enlarged cervical glands, 690 Health organization national, 166 Fractures, three unusual, 379 Freedom, medical, friends of, 388 Freire's, Dr., yellow fever inoculations, 335, 389, 502, 516
Friedenwald, cholesteatoma of the ear, 253
Frontal lobe, left, successful removal of tumor of, 16 State Boards of, conference of, 413 Heart-beat, self-regulation of, 573 successful operation for incised wound of, 300 tonic, a diuretic and, 384 Hematuria, renal, lasting twenty years, 692 GALL-BLADDER, extirpation of, 271 Hemoglobinuria, paroxysmal, 356 Hemorrhage, "accidental," with Gangrene of the great toe, 457
pulmonary, from the presence of a
tooth in a bronchus, 355 with uterine inertia, 159 Hemorrhoids, for, 18 symmetrical, in an infant, 573 painful, 190. Henry, angina pectoris, 615
Hepatic abscess, suppurative pylephlebitis and, secondary to appendicitis, 397
Heredity, inexorable facts of, 601 Gasserian ganglion, destruction of, for tri-geminal neuralgia, 183 Gaston, Dr. Freire's yellow fever inoculations, 335, 502
Gastric ulcer, etiology of, 34
Gastrostomy, esophageal stricture and, 294
for the removal of a peach-stone from Hernia, complications of, 454 diaphragmatic, 215 new operation for radical cure of, 104 of the bladder, 356
of the cecum in the inguinal canal,
eclampsia in children induced by, 626
of the placenta through the muscular the esophagus, 189 Giffen, cyst of pancreas; abdominal section; recovery, 626 Girls, young, treatment of painful prolapsed coat of the uterus during labor, 77 ovaries in, 12 Hernias, report of operations upon, 95
Herpes zoster, infectiousness of, 164
Herrick, report of seven cases of rupture of
the urinary bladder, with observations on Gladiolus, excision of, for sarcoma sterni, Glands, cystic degeneration of enlarged cervical, 690 diagnosis, 197
Hershey, prophylaxis of scarlet fever, 435
Herzog, cough of nasal origin, with report ductless, 497 Glaucoma in young persons, 438
Glycerin, neutral, therapeutic influence of subcutaneous injections of, 466
Goiter, exophthalmic, for, 244 a case, 687 Hessler, case of Norway itch, 517 Hiccough, for, 162 relieved by the wet-pack, 272 Himmelsbach, large doses of nitroglycerin, hemorrhagic, in the course of pregnancy, 132 reduction of, by the faradic current, 55 Gold and manganese, the use of in tuber-Hip-disease, pathologic classification of, culosis, 378
bichlorid of treatment of dipsomania, 403 Hirst, B. C., possible abuses in private hospitals, 54 Gold cures" in Colorado, 414 Holsten, report of a case of dermatitis medicamentosa from iodoform, potassium Gonorrhea of the rectum, 325 virulence of, 106 iodid, and aristol, 240 Homatropine susceptibility, 78 Goodell, remarks on a case of tubal pregsusceptibility to, 213 Homeopathy, end of, 577 in Germany, 472 one kind of, 600 nancy, 405 Gottstein's curet, a modification of, 382 Gould, a case of homatropine susceptibility, Hope, abscess of liver complicating cir-rhosis; multiple aspiration; recovery, 352 a temporary change in the axis of astigmatism, 598 the spelling of some medical words, 660 Gout, calomel for, 692 Graham, neuralgia at the base of the fourth Horse-leech as a human parasite, 553 Hospital, Anglo-American, at Rio Janeiro, in Philadelphia for infectious diseases, toe and its treatment, 512 Graves, excision of the gladiolus for sarcoma

purposes, State aid for, 248

out expense, 248

reports, a method of publishing, with-

Hospitals for infectious diseases, the right

to establish, 276 private, possible abuses in, 54 relief, at Columbian Exposition, 443

village and emergency, 533
Hubbard, report of the treatment of a case of fracture of femur during delivery, 323 Human being with a horse-mane, 498
Humerus, fracture of, during delivery, 697 Hyde, persistent stigmata resulting from cutaneous diseases, 281

Hydriatic or hydriatric, 28

Hydriatic or hydriatric, 28
Hydrocele, radical cure of, 494
Hydrogen dioxid, 193, 303
"Hydro-naphthol," the identity of, and impure beta-naphthol, 348

Hyperphoria, prominent symptoms of, as illustrated by thirteen successive cases,

Hysterectomy, Cesarean, in a malacosteon, 354 Hysteria, 101

Hysterical origin, choreoid movements in an adult male, probably of; relationship to habit-chorea, 231

ICE-COMMISSION, State, 437 Imbecility, moral, the opium-habit, and feigning, in which forgery is the offence committed, 29

Immigration law, the new, 276

necessity of restriction of, 21
Impetigo contagiosa, for, 148
Inaugural address, delivered before the
Medical Society of the State of New

York, 151 Infanticide in private lying-in houses, 332 Infectious diseases, hospital in Philadelphia

Inoculation, protective influence of, 466 Insane, gynecology among, 276 treatment of, outside of asylums, 259

Insanity in twins, 353
periodic, in which the exciting cause
appears to be the menstrual function,

538 Intestine, small, incarceration of, by the

vermiform appendix, 244 rupture of, caused by tapeworm, 243 Intubation, 48

Iodoform-amblyopia, 574 creasote and guaiacol, to deodorize, irritation from local use of, 13

Iron, the relative utility of the various preparations of, in the treatment of anemia and chlorosis, 356 Ischio-pubiotomy, 244 Itch, Norway, 517

JACOBSON, some recent cases of appendici-Jambul in the treatment of diabetes melli-

tus. 46 Jaw, excision of the lower, for malignant

disease, 383 lower, congenital ptosis, with synchronous movements of the affected lid

and, 409
Joint-disease, post-hemiplegic, 421
Journal, Daily medical, 554
Journalsm, medical, 653, 665

udges also, preliminary necessary, 389 qualifications

KEATING, aluminum wire pessaries (woolwrapped), 15 Keeley system, 167 Keeleyism, 498 and Keeley methods, 477 Keen, craniectomy and age, 476 Kidney, fibrinous coagula from, 520

intrinsic nerves of, 355 King, a case of placenta previa, with con-

clusions, 464
Kirkendall, conservative surgery in appen-

dicitis, 615 Klinedinst, a foreign body impacted in the larynx seventeen months; removal by laryngo-tracheotomy; recovery, 326

Knee-jerk, qualitative variations in, 691 Knowledge, popularization of medical and

hygienic, 221 Kreider, a case of movable or floating liver, 436

Kuh, a case of suffocative laryngo-tracheitis and a method of immediate relief, a topical treatment of bronchitis, 267

LABOR, complicated, 132, 328 delayed by rigidity of the os uteri, 54 hernia of placenta through muscular coat of uterus during, 77

premature, with symphysiotomy, 355 venesection in rigidity of the os uteri and in allied conditions that may retard, 305

Lactic acid, injections of, in the treatment of malignant growths, 18 amadrid, labor delayed by the rigidity of

the os uteri, 54
Lambert, relative localization of carcinoma in females, 306 Lancet relief fund, 166

Laryngitis, tuberculous, treatment of, with modified tuberculin, 88

Laryngo-tracheitis, suffocative, and method of immediate relief, 409

Larynx, a foreign body impacted in, seven-teen months; removal by laryngo-trache-

teen months; removal by laryngo-trache-otomy; recovery, 3:6

Lawrence, jambul in the treatment of dia-betes mellitus, 46

Lawrie, chloroform and the heart, 53

Lead-poisoning, chronic, elimination of

lead in, 396 Leffmann, analytic notes on liquid malt-extracts, 100

on some commercial disinfec-

tants, 395 Legislation, medical, in Georgia, 600 in Pennsylvania, 389 Leprosy in India, 552

Lesions, traumatic, of the spinal cord, 288

Leukemia and pseudo-leukemia, 191 Leukorrhea of children, for, 300 Life-insurance and the responsibility of the medical examiner, 358 tuberculosis and, 195 of the nervous American, duration of,

229 Liver, abscess of the, complicating cir-

rhosis, 352 movable or floating, 436 Live-stock industries, State bureaus of,

Lockridge, diabetes mellitus, following incised wound of the head, 28 Long, aids to the adoption of the metric system in prescription-writing, 322

Loomis, valedictory address, 74
Lowry, a case of "accidental" hemorrhage,
with uterine inertia, 159

Lupus erythematosus, 433 vulgaris, 433 Lusk, mitral stenosis in pregnancy, with

cases, 41 Lydston, a series of interesting cases of inflammation about the caput coli, 561 Lyman, epilepsy; cervico-brachial neuralgia,

Lynds, hernia of the placenta through the muscular coat of uterus during labor, 77 MAGAZINE, new medical, one method of launching, 553
Makuen, a modification of Gottstein's curet.

382 Malacosteon, Cesarean hysterectomy in, 354

Malaria, for, 384
Malarial origin, nephritis of, 694
Malignant degeneration in benign growths, 1 disease, excision of lower jaw for, 383 growths, lactic acid in treatment of, 18

new-formations, treatment of, 469
Malt-extracts, liquid, analytic notes on, 100
Manganese, the use of gold and, in tuberculosis, 378

Mania, acute, enormous doses of paralde-

hyde in a case of, 48

Manley, brachial palsy from fracture of
the clavicle; ischio-rectal abscess, 129 Masland, ovarian cyst with internal hemorrhage, 131

Massey, treatment of painful prolapsed ovaries in young girls, 12

Mastoid abscess, 203
Mattison, trional, the new hypnotic, 487
Mays, ice in the treatment of acute pneumonia, 681

croupous pneumonia, 68 IcCollom, J. H., precautions against Asiatic cholera in Boston, 601 McCollom, McGuire, the address of the President of the American Medical Association, 617

McKennan, symphysiotomy in private practice. 130

Measles, smallpox, scarlatina, and syphilis, etiology of, 410

with relapse, 216
Medical examiners bill, 525, 577
examining and licensing boards, State, conference of, 524 life, conduct of, 498

Medicine as a career, 304
"companies" and their methods of
doing business, 497

nobility of, 524 practice of, in Ontario, Canada, 528 practice of, the business aspect of, 600 Medicines, new method of administering, 384 patent, legal control of, 443

Memory or intellectual fitness, a question of, 524 unconscious, 413

Mercurial injections, pulmonary embolism from, 18

intoxication, 78
Merrill, the use of gold and manganese in tuberculosis, 378

Mesmerism, the new, 248 Methylene bichlorid, 217

Metric system, 420 aids to the adoption of, in prescription-writing, 322
practical helps in the use of, 362
Microscopic diagnosis, low-power obje

tives in, 26 Midwifery, antiseptic, 28 practice of, in New York, 555

Migraine, for, 550 Milk-diet, abuse of, 300 in nephritis, 494 inspection in Chicago, 578

legislation, 442 sterilization, 301
Mitchell, S., wood-sorrel as a cancer-cure,

363 S. Weir, choreoid movements in an

adult male, probably of hysterical origin; relationship to

habit-chorea, 231
hysteria; psychic anesthesia for
touch; psychic anosmia; psychic blindness in its variations, with comments, 101

post-hemiplegic pain; pre-hemi-plegic pain; post-hemiplegic sjoint-disease, post-hemiplegic nodes, 421

Mitral stenosis in pregnancy, 41 Montgomery, prolapsus uteri, 209 Moore, surgical interference in typhoid fever, 55 Moorehead, peculiar injury of eye, 573 Morton, a case of cholecystotomy, with re-covery after removal of gall-stones, 296 "Mountain fever," 376 Moyer, a rare occupation-neurosis, 188 Munn, diphtheria, a clinical study, 309, 337
Myelitis, traumatic, 235
Myringectomy, partial, and removal of incus
and stapes for relief of lesions of chronic

catarrhal otitis media, 509 Mythology, Norse, and cod-liver oil, 276 Myxedema, a new preparation of the thy-roid gland for treatment of, 276 terminating fatally in the course of thyroid treatment, 574 following bronchocele cured by thyroid therapy, 438

NEPHRITIS, milk-diet in, 494 of malarial origin, 694 without albuminuria, 391
Nerves, degeneration and regeneration of,

Neuralgia at the base of the fourth toe and its treatment, 512 cervico-brachial, 489 traumatic, relation of atmospheric elec-

tricity, magnetic storms, and weather elements to, 468 trigeminal, stretching of the facial nerve for the relief of, 695

Neuritis, puerperal, 17
Neurosis, traumatic, changes in the fields of vision in cases of, 78
New Way, 47
New York Academy of Medicine and Cro-

ton-water contamination, 303 Physicians' Mutual Aid Association, 360 Night-sweats of pulmonary tuberculosis,

for, 356 Nitro-glycerin for vomiting, 494

large doses of, 14 Nitrous oxid, chemistry and therapeutics of, 393

Noble, symphysiotomy versus its substi-tutes, with the report of a case of symphysiotomy, 176
Nodes, post-hemiplegic, 421
Noma in an adult, 190

Norbury, insanity in twins, 353 Northrup, Dr. W. P., 627 Nucleins, germicidal properties of, 536

MAJE

OBJECT-LESSON, 414
Obstetric cases, two, in which chloroform had no effect, 598 examination as taught by Credé and Leopold, some thoughts on, 250 Obstruction, intestinal, cholecystitis simulating, 383

Occupation-neurosis, rare, 188
Ointment-bases, 449
Ontario, Canada, practice of medicine in, 528 Oöphorectomy, secondary celiotomy fol-

lowing an, 542
Ophthalmia among hop-pickers, 600
Opium-habit, moral imbecility and feigning,
in which forgery is the offence committed, report of cases of, 29 Orbit, shot-wound of, 484 Orchiococcus, 442
"Orificial surgery," journal and devotees

of, 442 Orthopedics, plaster-of-Paris in, 70 Os uteri, labor delayed by rigidity of, 54 Ossicles, removal of the drum-head and, in diseases of the middle ear, 400

Osteoarthropathy, acute pulmonary, hyper-

trophic, 692
Osteomalacia during pregnancy, oöphorectomy for the relief of, 598 media, chronic catarrhal, partial myringectomy and removal of the incus and stapes for relief of lesions of, 509 Ovarian cyst with internal hemorrhage, 131

Ovaries, painful prolapsed, in young girls, treatment of, 12

PACKARD, J. H., traumatic separation of lower epiphysis of femur, 26 Page and Hutchinson, severe fracture of the vertex, with replacement of a large fragment of bone, 185

Pain and shock, 666

Pain, post-hemiplegic, 421
pre-hemiplegic, 421
Palsy, brachial, from fracture of the clavi-

Pan-American Medical Congress, Germany and, 247 official languages at, 165

publication of papers read at, 221

Pancreas, cyst of, 626 Papers, exclusive publication of, 443

limit, to ten minutes, 303
Paraldehyde, enormous doses of, in a case of acute mania, 48 for spasmodic asthma, 187

Paralysis, acute ascending, fatal case of, 161 agitans, sodium borate in the treatment

of. 216

for trigeminal neuralgia, with a re-port of two cases, 183 esophageal stricture and gastrostomy,

294
Parker, interesting manifestations of syphilis in the eye and throat; ulcers of the epiglottis, strabismus, and paralysis of ocular

glottis, strabismus, and paralysis of ocular muscles, 463
Parotiditis, complications of, 443
Parsons, a contribution to the subject of acute pleurisy; its pathology, etiology, symptomatology, and treatment, 57
Patella, fracture of, treated by continuous extension; patients not confined to bed, 5
Pathology and biology, 360
Peckham, study of a case of erysipelas genitalium, due to the use of infected ointment 148

ment, 148

Pelvis, unique fracture of, 17 Pensions for physicians' widows, 627
Pepper, two cases of pulmonary tuberculosis: one chronic, one acute, 544
Perineum, to prevent laceration of, by shoulder, 12
Pessaries, aluminum wire (wool-wrapped),

Peterson, a case of appendicitis occurring on the seventh day following labor; rupture into the bowel; recovery, 515 the treatment of insane outside of asy-

lums, 259
Pharmacist, physician and, 277
Pharmacists, physicians as, 194, 278
Philanthropy, munificent, 525
Phlegmons, emphysematous, etiology of, 438 Physician, a true, 51 and pharmacist, 277 and the sectarian, 220 Physicians as pharmacists, 194, 278

hospital, unmerited censure of, 523 hotel, again, 414 legacies to, 304
Physio-medical grammar and therapeutics,

Pilcher, inaugural address, delivered before the Medical Society of the State of New

York, 151 Placenta, hernia of, through the muscular coat of the uterus during labor, 77

Placenta previa, 464
Plaster-of-Paris in orthopedics, 70
Platt. obstacles to a national quarantine,

Pleurisy, a contribution to the subject of acute, 57 Plumbism, for painful constipation of, 574

Pneumonectomy, 471
Pneumonia, acute, ice in the treatment of, 68x

apical, 549 bilateral polyneuritis following, 438 croupous, ice in the treatment of, 68 Pneumotomy twice in the same patient for the relief of tuberculous abscess and gangrene of the lung; recovery, 38 Pohlman, duration of life of the nervous

American, 229
Poisoning by sardines, 216
deaths from, in England and Wales, 275

"Politics," 305
American, as affecting public medical officers, 473

Polymastia, 464
Polyneuritis, bilateral, following pneumonia, 438

Poor, model dwellings for, 136 Porcher, wood-sorrel for carcinoma, 363 Porteous, eclampsia gravidarum et parturientium, 145
Posterior root, the relation of, to the pos-terior horn in the medulla and cord, 189

Practice, medical, in Connecticut, regula-

rractice, medical, in Connecticut, regula-tion of, 248 in Ohio, regulation of, 362 of medicine in Connecticut, 442 in Ontario, Canada, 528 Precocity extraordinary, 333 Pregnancy, hemorrhagic goiter in the

course of, 132
incoercible vomiting and salivation of, requiring abortion, 162
mitral stenosis in, 41 precocious, 574 tubal, remarks on a case of, 405 Preston, traumatic lesions of spinal cord,

Prince, obstruction of bowel from Meckel's

diverticulum; a cholecystotomy, 45 Profession, English, medical statistics of, 414

medical, as a calling, Dr. Billings's estimate of, 221 of Cincinnati, Ohio, 333 salaries of, 304 the, and public scandal, 137

Prolapsus uteri, 209
Proprietary preparations, taxation of secret, 471

Prostitution, suppression of, 414

Prurigo, 162
Pseudo-leukemia, leukemia and, 191
Pseudo-leukemia, leukemia and, 191
Ptosis, congenital, with synchronous movements of the affected lid and the lower jaw, 409

Pubeotomies, two, 47 Pubio-ischiotomy, unilateral, operation of,

249 Puff-ball as a hemostatic, 190 Pulmonary artery, prenatal closure of, 437 gangrene from the presence of a tooth in a bronchus, 355

Purgatives, the effect upon sucklings of, administered to the mother, 356

Purpura hemorrhagica, 10
simplex, infectious nature of, 493
Pylephlebitis, suppurative, and hepatic abscess secondary to appendicitis, 397
Pyloric obstruction, 263
Pyloric obstruction, 263 Pyoktanin for ptyalism, 356

QUACKS, tax on, 22 preliminary, necessary, Qualifications, judges also, 389

Quarantine and sanitation at the port of Philadelphia, legislation in reference to, 473 at Hamburg, 471

bill, Senate, 79 committee of the New York Academy of Medicine, 247

fin-de-siècle, 472 national, obstacles to, 27 Quinine for cholera, 520 for chorea, 466

RACHFORD, the etiology of chorea, 429
Rags, infected, importation of, 303
Randall, epilepsy or angina pectoris, 277
Rectum, gonorrhea of, 325
Reformatory, New York State, 554
Reichert, a comparative study of the physical siologic actions of brucine and strych-

nine, 369 Remedy, the single, 525

Review

Adler, Fissure of the Anus and Fistula

in Ano, 138
Berry, Diseases of the Eye, 528
Beyer, First Aid to the Injured and
Transportation of the Wounded, 280 Transportation of the Wounded, 280
Borcicke and Dewey, The Twelve Tissue Remedies of Schüssler, Comprising the Theory, Therapeutic
Application, Materia Medica, and
a Complete Repertory of these
Remedies, Homeopathically and
Bio-chemically Considered, 695
Braithwaite, The Retrospect of Medicine 276

icine, 556

Brockway and O'Malley, Anatomy, 108 Cerna, Notes on the Newer Remedies,

Dexter, The Anatomy of the Perito-

neum, 696
Ewald, Diseases of the Stomach, 279
Foster, An Illustrated Encyclopedic Medical Dictionary, 23

Galton, Finger-prints, 279 Gowers, Syphilis and the Nervous Sys-

tem, 527 Gray, Treatise on Nervous and Mental

Diseases, 415 Grünwald, Die Lehre von den Naseneiterungen. u. s. w. (The Suppurative Processes in the Nasal Passages, with Special Reference to the Diseases of the Ethmoid and Sphe-

noid Bones), 138 Hansell and Bell, A Manual of Clinical Ophthalmology, 139 Hirst and Piersol, Human Monstrosi-

rins and Piersol, riuman Monstrosi-ties, Part IV, 526
Ingals, Diseases of the Chest, Throat, and Nasal Cavities, 83
Jackson, Ready Reference Handbook of Diseases of the Skin, 23
Kenner, Contributions of Physicians

to English and American Literature,

Luff, A Manual of Chemistry, 278 Lydston, Gonorrhea and Urethritis, 418 Manning, Physiology: A Manual for Students and Practitioners, 696 Marcy, The Anatomy and Surgical

Treatment of Hernia, 415
Mathews, A Treatise on Diseases of the Rectum, Anus, and Sigmoid

Flexure, 417 McLaughlin, Fermentation, Infection, and Immunity, 418
Miller, Die Microörganismen der
Mund-höhle, 418

Miller, McEvoy, and Weeks, Diseases of the Eye, Ear, Throat, and Nose,

Reviews-

Minot, Human Embryology, 25 Morris, Human Anatomy, 416 Morrow, A System of Genito urinary Diseases, Syphilology, and Dermatology, 526 Nuttall, Hygienic Measures in Rela-

tion to Infectious Diseases. non to Infectious Diseases, 25
Parkes, Hygiene and Public Health, 24
Pellew, A Manual of Practical and
Medical Physiological Chemistry, 555
Senn, Tuberculosis of Bones and

Joints, 24 Shoemaker, A Practical Treatise on Diseases of the Skin, 109

Stearns, Lectures on Mental Diseases, Designed Especially for Medical Students and General Practitioners,

Sternberg, A Manual of Bacteriology,

Swanzy, A Handbook of D.seases of the Eye and their Treatment, 139 Tanner, Memoranda on Poisons, 280

Tiffany, Tablets for the Measurement and Recording of the Magnified Image of the Fundus of the Eye,

Transactions of the American Gvnecological Society, 1892, Vol. XVII,

Transactions of the American Surgical Association, Vol. X, 527
Transactions of the Association of

American Physicians for the Year 1802, 270

sactions of the Colorado State Medical Society, 556
Transactions of the Medical Society of

the State of North Carolina, 556 Transactions of the Texas State Med-

ical Association, 1892, 139
Tuke, A Dictionary of Psychological
Medicine, 83
Wright and Northrup, The Diseases of

Children, Medical and Surgical, 697 Year-book of Treatment for 1893, 556 Rheumatism, articular, the employment of sodium salicylate by enema in the treatment of, 550

Rhinitis, fibrinous, etiology of, 536 membranous, e'iology of, 505

Rhizodontrophy, 359 Rochester, the metric system, 420 Rohé, sanitary condition of Baltimore, 419 Ross, to prevent laceration of perineum by the shoulder, 12

Roussel, a case of malignant syphilis resulting in death, 540 Rubber, dangers of, taken into mouth, 384

SALOL subcutaneously in the treatment of pulmonary tuberculosis, 132 Sangree, tic douloureux from visual defect, 572 Sanitarium Association of Philadelphia, 496

Sanitary condition of Baltimore, 419 conference, Dresden, 523 legislation in Pennsylvania, 663

precautions, 445 Sanitation in Philadelphia, 501

in Roumania, 600 quarantine and, at the Port of Philadelphia, legislation in reference to.

473 Sarcoma sterni, excision of gladiolus for, 241 Sardines, poisoning by, 216

Scabies, for, 244 Scandal, public, the profession and, 137 Scarlatina, bovine, 359 measles, smallpox, and syphilis, eti-

ology of, 410 Scarlet fever, prophylaxis of, 435 Scheming and frankness, 137

Schemm, a case of aneurism of the aorta associated with insufficiency of the aortic

valves, 297 Scheppegrell, an improved vapor-appara-

tus, 353 Schmid, some personal observations and reflections upon alcoholism, the effects of alcoholic abuse upon posterity, and the treatment of alcoholism, 90 Schneider, influence of diseases of the ear

upon the mental and physical development of the child, 372

Scholarships to medical students, 577 Schools, sanitary supervision of, 304 Schroeder, mastoid abscess, 203 Schulin, a combination of dropper and

hypodermatic syringe, with improve-ments on both, 518

ciatica, primary, 691 Sclerosis, association of posterior spinal, and general paresis, 78. posterior spinal, early eye-symptoms

of, 355 Sectarian, the physician and, 220 Sectarianism, medical, medical students,

and, 195 Senility, dysuria of, 412 Sewage, disposal of, 8

Shattuck, some remarks on arsenical pois-Shattuck, some remarks on arsenical pus-oning, with special reference to its do-mestic sources, 589 Shimwell, the reason why the neck of the femur does not break, 97

Shock and pain, 666
Shoemaker, G, E., recovery after swallowing
a teaspoonful of amyl nitrite, 544
annuality pylephlebitis and suppurative pylephlebitis and hepatic abscess secondary to appendicitis, with a report of

two cases, 397
J. V., ointment bases, 449
Shoulder-joint, left, subcoracoid dislocation of, 457

Simpson, physician and pharmacist, 277 Singultus, for, 410 hysterical, 127

Skeel, the use of the curet in abortion, 207 Sleep, a new theory of, 161 seizures, periodic, epileptic, 598 Smallpox at Leicester, England, recent out-

break of, 414
epidemic at Antwerp, 554
measles, scarlatina, and syphilis, eti-

ology of, 410 Smith, shot-wound of the orbit, 484

Sneve, an operation for re-covering the denuded cranium, 239 Society Proceedings

American Academy of Medicine, 677, 702 American Association of Obstetricians

and Gynecologists, 640 American Climatological Association,

American Gynecological Society, 558, 578

American Laryngological Association, 603 American Medical Association, 643,

667, 697 American Pediatric Society, 637

American Surgical Association, 614, 631 Association of American Anatomists, 109 Association of American Physicians.

612, 628 Medical Society of the State of Penn-

sylvania, 556, 587 New York Neurological Society, 221,

333, 445
Philadelphia County Medical Society,

Twenty-second Congress of the Association of German Surgeons, 499 Solly, neglect of the early diagnosis and treatment of pulmonary tuberculosis, 117

Soya beans in diabetes mellitus, use of, 662

Speech without a larynx and without use

Tapeworm, rupture of the small intestine caused by, 243

Tariff, protective, and the profession, 303

Typewriter for the blind, 360

Typhold and cholera excreta, disposal of, 359 of the lungs, 195 Spelling of some medical words, 660 Teacher, the medical, 245
Terpene hydrate for bronchial catarrh, 384 Testimony, expert, 467 medical, 627 reform, 665 Spine, curvature of, lateral, new aid in the diagnosis of, 452 "Spitting" habit, 411 Tetanus, preventive treatment of, 110 Tetronal, trional and, 494 Teucrin, a new therapeutic agent, 272 Splenectomy, successful, for wandering spleen, 244
Spoils, division of, 359
Sponges for surgical purposes, 244
Sporozoa and their relation to carginoma, Therapeutic notes, 351, 596
Therapeutics, biologic, 215, 551
Therapy, animal products in, 246
Thigh, two rare tumors of, 271 Thompson, two obstetric cases in which chloroform had no effect, 598

Thorner, treatment of tuberculous laryn-Stamm, report of a case of a large abdom inal echinococcus cyst, with remarks, 365 Starr, craniectomy and age, 531 State Bureaus of Live-stock Industries, gitis with modified tuberculin, 88 Thyroid feeding, sporadic cretinism suc cessfully treated by, 662 Statistics, illustrations of the fallacies of, gland, the function of, and the therapy of thyroid disease, 165 therapy, myxedema following broncho-166 vital, concealment of, 388 Steele, plaster-of-Paris in orthopedics, 70 Steele, plaster-of-Paris in orthopedics, 70
Stelwagon, syphilis, epithelioma, lupus
vulgaris, and lupus erythematosus, 433
Stenography, medical, the business of, 666
Stenosis, mitral, in pregnancy, 41
Sternberg, Dr. Freire's yellow-fever inoculations, 380, 616
George M., 627
Stewart, a résumé of some modern methods cele cured by, 438
treatment, a case of myxedema that
terminated fatally in course of, 574
Tie douloureux from visual defect, 572 Time allowed speakers and essayists, 414
Tobacco, alcoholic liquors and, should they
be furnished to prisoners? 443 Tolypyrin, 410 Tolysal, 410 of the diagnosis of the diseases of the of the diagnosis of stomach, 169 rephritis without albuminuria, 391 and Beckwith, the identity of the pro-so-called "hydro-naph-Tomes, a new aid in the diagnosis of lateral curvature of the spine, 452
Tongue, black, 275
position of, in peripheral palsy of facial prietary, so-called "hydro-naph-thol" and impure beta-naphthol, 348 nerve, 18 Stigmata, persistent, resulting from cuta-neous disease, 281 Stockton, hysterical singultus, 127 Tonsillitis, acute, 461 Tonsillotome and uvulatome, combined, the etiology of gastric ulcer, 34
Stomach, a résumé of some modern methods of the diagnosis of diseases of, 169 Trachea, profound opening of, for the re-moval of a foreign body, 573 Tracheotomy in an infant three days old, Strabismus, non-operative treatment of, 497 Street-sweepings and rubbish, 498 243 Tremaine, amputation of lower part of Tremane, amputation thigh, 548
Trichinæ in hogs, 333
Trional and tetronal, 494
the new hypnotic, 487
Tyron, Surgeon-General of the Navy, 554
Tuberculin, modified, some experiments Stricture, esophageal and gastrostomy, 294 Strychnine, brucine and, a comparative study of the physiologic actions of, 369 Students, medical, and medical sectarianism, 195 Study, more years of, or more study in the year? 358
Substitution, 499
Surgical, medical and, exhibit, 389 with, 85 treatment of tuberculous laryngitis with, 88 Sycosis, for, 190 Tuberculosis and life insurance, 195 Symphysiotomies, two, one for the extracfor, 692 gold and manganese in, 378 infectiousness of, 471 tion of a decapitated and imprisoned head, 271 Symphysiotomy, 328 of the bladder, 457 predisposition to, 132
pulmonary, creasote treatment of, 132
for the night-sweats of, 356
neglect of early diagnosis and a case of, 272 first, in London, 249 in a male, 300 in private practice, 129 premature labor with, 355 treatment of, 117 versus its substitutes, with the report of a case of symphysiotomy, 176 salol subcutaneously in the treatment of, 132 two cases, one chronic, one acute, Syphilis, 433 hereditary, 544 venesection in, 246 hind, 433 hereditary, a hitherto undescribed symptom of, 328 in the eye and throat, 463 Tuberculous abscess and gangrene of the lung; pneumotomy twice in the same malignant, resulting in death, 540 patient, 38 and typhoid infection, mixed, 219 measles, smallpox, scarlatina and, etiinspiration-broncho-pneumonia, 210 larnygitis, treatment of, with modified tuberculin, 88 ology of, 410 treatment of, 156
by means of hypodermatic injections of mercury, 550
Syringe, hypodermatic, combination of dropper and, with improvements on both, meat, the proportion of, 34r the use of, 348 patients, need of special hospitals for treatment of, 166 tissues, treatment of suppurating, 692 Tuholske, pyloric obstruction, 263
Tumor of left frontal lobe, successful re-

TALIPES valgus and talipes calcaneus, 269 Tammany versus the people of the United States, 166

moval of, 16

Tumors, two rare, of the thigh, 271

fever, relapses of, 154 surgical interference in, 55 water-supply and, 442 infection, mixed tuberculous and, 219 Typhus fever in New York, 166 ULCER, gastric, etiology of, 54 of the stomach, 121 University of Pennsylvania, professorship of University of remarkation, pro-mental diseases, etc., 524 Uremia in an alcoholic patient with visceral complications successfully treated by venesection, diaphoresis, and nitroglycerin, 298 Ureter, resection of, 17 Urethritis, a new instrument for the treatment of, 15 Urinalysis, some notes upon, 201 Urine, a gas-generating bacterium in, 190 Uterus, a new method for rapidly removing, and its appendages, antiseptic cata-phoresis in the treatment of diseases of, 65 hernia of the placenta through the muscular coat of, during labor, 77
rupture of, during labor, with the
placenta in the abdominal cavity, 691 severance of cervix from body of, 243 spontaneous closure of rupture of, 162 Uvulatome, tonsillotome and, combined, 14 VALEDICTORY address, 74 VALEDICTORY address, 74
Vapor-apparatus, an improved, 353
Vaughan, Novy, and McClintock, the
germicidal properties of nucleins, 536
Veasey, experiments with cocaine phenate
as a local anesthetic, 345
Veeder, reduction of goiter by the faradic current, 55 Venesection, 303 in pulmonary tuberculosis, 246 in rigidity of os uteri and in allied conditions that may retard labor, 305 in thoracic aneurism, 520 Vermiform appendix, removal of, III Vertex, severe fracture of, with replacement of a large fragment of bone, 185 of a large fragment of bone, 185
Vertigo, aural, in children, 409
Vienna, University of, 498, 577
new buildings for the surgical
clinics of, 249
Viscera, transposition of, 598
Vision, changes in the fields of, in cases of traumatic neurosis, 78 Vomiting, nitroglycerin for 494 WALKER, cancer-cure, 110
Warthin, additional notes on the diazoreaction, 569 Washburn, albuminuria in the apparently healthy, 341 Water, additional bacillus found in, 244 Water, additional bacillus found in, 244
filtration, etc., at Hamburg, 444
supply and typhoid fever, 442
cholera and, 332
of Philadelphia, 441
Wents, abortive free crucial incision in the
treatment of carbuncle, 307
Wet-pack, hiccough relieved by, 272
Whelpley, physicians as pharmacists, 278
White, pneumotomy twice in the same patient for the relief of tuberculous abscess
and gengrane of the lung; 280 who oping-cough, etiology of, 104 Wilard, complications of hernia, 454 talipes valgus and talipes calcaneus, 269

Williams, E. H., low-power objectives in microscopic diagnosis, 26
Talcott, pain and shock, 666
J. W., a new instrument for the treatment of urethritis, 15
Wire-suture twister, an improved antiseptic, 354
Wire-suture twister, an improved antiseptic, 354
Woodward, a case of transposition of viscera, 598

Work, "mountain fever," 376

YELLOW-FEVER inoculation, 335, 502 inoculations, Dr. Freire's, 389, 616 Young, a pathologic classification of hip-disease, 403

A WEEKLY MEDICAL JOURNAL.

ol. LXII. No. 25.

SATURDAY, JUNE 24, 1893.

Per Annum, \$4.00. Per Copy, 10 Cents.

Contents.

ORIGINAL ARTICLES:

Ice in the Treatment of Acute Pasumonta. By TEOMAS J. MAYS, M.D., of Philadelphia Cough of Nasal Origin, with Report of a Case. By MAXIMILIAN HERZOG, M.D., of Cincinnati, Ohio 687

CLINICAL MEMORANDA.

Traumatic Empyema Communicating with the Bowel. By N. T. DULANEY, Cystic Degeneration of Enlarged Submaxillary Glands. By D. W. Havs, M.D., of Osawatomie, Kan 690

MEDICAL PROGRESS

A Case of Diphtheric Croup in which a Tracheotomy-tube was Worn for Sixty Days-Rupture of the Uterus during Labor, with the Placents in the Abdominal Cavity-Qualitative Variations in the Knee-jerk-Primary Sciatica 691 Renal Hematuria, Lasting Twenty Years-Acute Pulmonary Hypertrophic Osteoarthropathy 692

THERAPEUTIC NOTES.

The Treatment of Suppurating Tuberculou Tissues - Bromoform-intoxication - For Tuberculosis-Calomel for Gout-To Allay the Craving for Alcohol 692

EDITORIAL ARTICLES.

The Report of the Committee on Revision of the Code 693 Nephritis of Malarial Origin . . . 694

EDITORIAL COMMENTS.

Stretching of the Facial Nerve for the Relief of Trigeminal Neuralgia-The Etiology of Friedreich's Ataxia-Professional Confidence-Bacillus Aërogenes Meningitidis-Ecphyaditis 695

REVIEWS.

The Twelve Tissue Remedies of Schussler. By William Boericke, M.D., and Willis A. Dewey, M.D. 695 Lectures on Montal Diseases. By Henry Putnam Stearns, A.M., M.D. 696 The Anatomy of the Peritoneum. By Franklin Dexter, M D. 696 Physiology, A Manual for Students and Practitioners. By Frederick A. Manning, M.D. (The Students' Quiz Series) 696 Transactions of the American Gyne-

Tablets for the Measurement and Recording of the Magnified Image of the Fundus of the Eye. By Flavel B Tiffany, M.D. 697 ases of Children, Medical and Surgical. By Henry Ashby, M.D Lond., F.R C.P., and G. A. Wright, F.R.C.S. . 697

CORRESPONDENCE.

Fracture of the Humerus during De-

SOCIETY PROCEEDINGS.

American Medical Association. Fortyfourth Annual Meeting, held at Milwaukee, Wis., June 6, 7, 8, and 9, 1893. Section on Diseases of Children. First day, June 6th 697 Second day, June 7th 699 Third day, June 8th 700 American Academy of Medicine. Righteenth Annual Meeting, held at Milwaukee, Wis., June 3 and 5, 1893.

NEWS ITEMS.

Fatal Quackery-Pasteur Institute in the Punjab - Paul Guttmann-James Mole-Books and Pamphlets Received 704

Entered at the Post Office at Philadelphia as Second-Class Matter. Copyright 1893, by Lea Brothers & Co.

Smith on Children.—Seventh Edition.

A Treatise on the Diseases of Infancy and Childhood. By J. Lewis Smith, M. D., Clinical Professor of Diseases of Children in the Bellevue Hospital Medical College, New York. Seventh edition, thoroughly revised and rewritten. In one handsome octavo volume of 881 pages, with 51 illustrations. Cloth, \$4.50; leather, \$5.50.

Seiler on the Throat and Nose.—New (4th) Ed. Just Ready.

A Handbook of Diagnosis and Treatment of Diseases of the Throat, Nose and Naso-Pharynx. By CARL SEILER, M. D., Lecturer on Laryngoscopy in the University of Pennsylvania. New (fourth) edition. In one handsome 12mo, volume of 214 pages, with 107 illustrations and 2 colored plates. Cloth, \$2.25.

Taylor's Medical Jurisprudence. New Edition. Just Ready.

A Manual of Medical Jurisprudence. By Alfred S, Taylor, M. D., Lecturer on Medical Jurisprudence and Chemistry in Guy's Hospital, London, New American from the twelfth English edition. Thoroughly revised by Clark Bell, Esq., of the New York Bar. In one octavo volume of 787 pages, with 36 illustrations. Cloth, \$4.50; leather, \$5.50.

Stimson on Fractures and Dislocations.

A Treatise on Fractures and Dislocations. By Lewis A. Stimson, M. D. In two handsome octavo volumes. Vol. I., FRACTURES, 582 pages, 360 illustrations. Vol. II., DISLOCATIONS, 540 pages, with 163 illustrations. Complete work, cloth, \$5.50; leather, \$7.50. Either volume separately, cloth, \$3.00; leather, \$4.00.

Lea Brothers & Co., Publishers, 706, 708 & 710 Sansom Street, Philadelphia.

URIC ACID DIATHESIS

Treatment with Buffalo Lithia Water

By eminent physicians as follows

DR. HUNTER MCGUIRE, says:—"In Uric Acid Gravel, and indeed, in diseases generally dependent upon an Uric Acid Diathesis, Buffalo Lithia Water Spring, No. 2, is a remedy of extraordinary potency. I have prescribed it in cases of Rheumatic Gout, which had resisted the ordinary remedies, with wonderfully good results. I have used it also in my own case, being a great sufferer from this malady, and have derived more benefit from it than from any other remedy."

DR. J. T. LeBLANCHARD, Professor Montreal Clinic, SM., SN., V. U. "I have used Buffalo Lithia Water in the most obstinate cases of Chronic Inflammation of the Bladder, in Stone of the Bladder, in Uric Acid Gravel, with the most efficacious results."

DR. WM. B. TOWLES, Professor of Anatomy and Materia Medica in the Medical Department of the University of Virginia. "Buffalo Lithia Water Springs, No. 2, has proved far more efficacious in many diseased conditions than any of the simple Alkaline waters. I feel no hesitancy whatever in saying that in Gout, Rheumatic Gout, Rheumatism, Stone in the Bladder, and in all diseases of Uric Acid Diathesis, I know of no remedy at all comparable to it. Its effects are marked in causing a disappearance of Albumen from the Urine."

DR. HARVEY L. BYRD, Of Baltimore, President and Professor of Obstetrics and Diseases of Women and Children in the Baltimore Medical College, formerly Professor of Practical Medicine, etc. "I have witnessed the best results from the action of the Buffalo Lithia Water, Spring No. 2, in Chronic Gout, Rheumatic Gout, Rheumatism, Gravel and Stone in the Bladder, and I do not hesitate to express the opinion that in all deseases depending upon or having their origin in Uric Acid Diathesis, it is unsurpassed, if indeed it is equaled, by any water thus far known to the Profession."

DR. ALLARD MEMMINGER, Professor of Chemistry, Medical College, State of South Carolina. "I have used in my own case and prescribed for others Buffalo Lithia Water for Uric Acid trouble with excellent results, and I regard it as the safest, surest, and most agreeable way of removing from the system this most pernicious derivative of Urea, Uric Acid, the retention of which is followed by so many distressing symtoms, embracing Gout, Calculi of the Kidney and Bladder, Herpetiform Neuralgic affections, cases of Mental Depression and Nervous Irritability and Nervous Asthma caused by the irritating action of Uric Acid on the bronchial tubes."

JAMES L. CABELL, A. M., M. D., LL. D., Professor of Physiology and Surgery in the Medical Department in the University of Virginia, President of the National Board of Health, etc., etc. "The Buffalo Lithia Water Spring, No. 2, contains in notable quantities two of the alkalies which are accredited as extremely valuable in the treatment of Gout, Lithiasis and Liver Affections. I refer to the Carbonates of Potash and Lithia. It is now well known that both of these alkaline carbonates have an ascertained value in cases of Uric Acid Diathesis connected with gravel, and in cases of Chronic Gout, because of their affinity for Uric Acid, and the great solubility of the salts which are formed by their union with that acid. If this were all, the range of their therapeutical indications would be comparatively restricted; for, though dissolving the Uric Acid already formed, they would tail to meet the most important indication—that of rooting out the diathesis on which the formation depends, It is, however, held by eminent medical authorities, that 'the beneficial effects of alkalies are not due to their neutralizing acidity, or to any direct action upon Uric Acid,' but that they 'seem to do good by combatting the pathological state on which the formation of Uric Acid depends." "We are not, however, left to theory, as to the action of this water, as numerous testimonials from the profession seem to have established the fact that it has already accomplished results, such as its composition would indicate." "In my experience it is a remedy of decided efficiency in the Lithaemic condition. I have also found it a direct tonic for the nervous system in cases of Cerebral Exhaustion. The nervous symptoms, however, in these cases may have been due to a Lithaemic condition."

Water in cases of one dosen half-gallon bottles, \$5.00, f. o. b. here. For sale by all first-class druggists

THOMAS F. GOODE, PROPRIETOR



OTHER

Farbenfabriken

PRODUCTS

SALOPHEN

A safe and effective derivative of phenol and salicylic acid. Indicated in rheumatism and allied maladies.

SULFONAL-BAYER

THE greatest of hypnotics for general uses, and invaluable in the neuroses. Flust be given as directed.

ARISTOL

A Prompt and Effective dressing, for all external and cavital lesions. Safe, convenient and odorless.

PIPERAZINE-BAYER

URIC Acid Solvent, and of remarkable efficacy in the uric diathesis. Indicated also in gout and renal colic.

TRIONAL

A hypnotic and sedative which is specially indicated in many neuroses not amenable to other drugs.

LOSOPHAN

A^N antimycotic, indicated in ringworm, scables, pityriasis, and the parasitical dermatoses in general.

EUROPHEN

SUBSTITUTE for iodoform, cicatrisant, and local dressing for catarrhal, ulcerative and luetic lesions.


EUROPHEN-ARISTOL

This is a mixture of these remedies. It is thought to combine the advantages of both.

Pamphlets forwarded by

W. H. Schieffelin & Co., New York.

Sole Agents and Licensees for the United States.



Combined with Vaseline or other vehicle, and used with this instrument, are Camphor, Carbolic Acid, Hamamelis, Zinc Oxide, Oil Eucalyptus, Menthol and other remedies.

The Treatment of Inflammation of the Mose and Throat in this manner is now universally regarded as the best.

Has screw neck vial.

Metal screw cap and tubes.

Handsomely nickel-plated.

Bulb of best rubber.

Easily taken apart for filling or cleaning.
No danger of spilling the

PRICE, \$1.00 NET. POSTAGE, 7 CENTS.

New Hinstrated Catalogue of Surgical Instruments, 300 pp., 2000 new illus., sent on receipt of 14s. in stamps for the nestage. New Pampalet on Atomization of Liquids mailed free on request.

DR. CHADWICK'S GYNECOLOGICAL TABLE.



This table is offered as a substitute for the unsightly, often ill-adapted and expen sive chairs commonly used

by gynecologists.
It is constructed of thoroughly-seasoned oak, well fitted and finished, and when not in use will pass for an ordinary desk. It is provided with hair-stuffed and hair-cloth covered mattress. leg and foot rests, making it adapted for Sims's position, dorsal position, etc., also with drawers and compart-

ments for instruments, tampons, pessaries, absorbents, and with glass-stoppered bottles and pots for acids, lubricants, etc.

The following advantages are claimed for this table:

- I. That its presence in an office will never excite suspicion.
- That it is perfectly adapted for examination in all positions.
- That in it may be kept readily accessible all the instruments, appliances and medicaments likely to be needed.

That its moderate price makes it easily obtainable by every physician.

Perfectly adapted to examinations and treatment on diseases of the rectum and genito-urinary organs, while often convenient as a general operating table.

Price, \$50.00. Boxing, \$2.50.

Circulars, with full descriptions and other cuts, sent on application. Many unsolicited commendatory letters, from physicians using this table, have been received by us.

CODMAN & SHURTLEFF, SUPERIOR SURGICAL INSTRUMENTS, 13 & 15 TREMONT STREET,

FEICK BROTHERS, Manufacturers of SURGICAL INSTRUMENTS, ORTHOPÆDIC APPARATUS, Etc.

No. 21 SIXTH STREET, PITTSBURGH, PA.

OUR COMPACT NEW OPERATING CASE No. 11.



CONTAINS:

I Amputating Knife, Leg and Arm, I Finger Knife, I Hernia Knife, I Sharp Curved Bistoury, 2 Scalpels, I Tenotome, I Tenaculum, I Pair Scissors, curved, I Finger Saw, I Capital Saw, I Liston's Bone Forceps, I Artery and Needle Forceps combined, I Dressing Forceps, I Esmarch's Tourniquet, with chain, I Director with Aneurism Needle, I Pair Probes, I Tablet Silk, four sizes, in slip case, I Coil Silver Wire, one-half dozen Assorted Needles.

In Fine Mahogany Case, lined with leather In Fine Morocco Leather Case, lined with leather -22 00 In Fine Mahogany Case, Instruments fitted in polished

cedar wood, no lining whatever, knives fitted in metalrak to lift out. The Finest Aseptic Case in the market

Any of the above with Aseptic Trephine and Trephine, Elevator in addition

All the instruments have our All Metal Antiseptic Handles. These are the only handles that can be boiled indefinitely without damage. They are hollow, therefore light; are of German Silver, hence susceptible of a high polish; are grooved to give a firm grasp; are soldered to the blades, consequently cannot become loose, and have no cracks or crannies to collect impurities.

Our new Illustrated Catalogue of Surgical Instru-ments, Deformity Apparatus and Artificial Limbs, containing 328 pages and over 2000 Illustrations, will be sent on application.

LARGER CASES HADE UP TO ORDER, AT MODERATE COST.

CARDS.

Cards occupying this space are inserted once for \$2.50, twice for \$4.00 and three times for \$6.00. Cash must accompany the order. Address ADVERTISING DEPARTMENT,

THE MEDICAL NEWS, Philadelphia.

Association of Hospital Physicians and Surgeons of Philadelphia.

Physicians visiting Philadelphia who desire to see clinical work in the Hospitals, can receive a Roster by calling upon the Secretary of the Association,

DR. H. R. WHARTON, 112 S. 18th St.

23 50

INTERNATIONAL PROICAL CONGRESS IN ROME

North German Lloyd S. S. Co.

EXPRESS STEAMERS TO GENOA

FOR PHYSICIANS ATTENDING

MEDICAL CONGRESS IN ROME.

Werra, Aug. 5:

Fulda, Aug. 19:

Werra, Sept. 9:

RETURNING FROM GENOA.

Kaiser, Oct. 4: Fulda. Oct. 11:

OR RETURNING FROM BREMEN EVERY SATURDAY.

For further particulars apply to

OELRICHS & CO.

2 BOWLING GREEN.

NEW YORK.

COLOR IN CALISAVA.

It is a remarkable fact that the alkaloidal principles of plants are perfectly colorless-are pure white. This is the more curious when we consider the variety and intensity of color in the vegetable kingdom.

Take, for example, some of our most important and best known drugs: the coca with its peculiar greenish color; the dark brown of the onium: the red and vellow calisava; vet their respective alkaloids or active principles-cocaine, morphine, quinine, etc., are all white. The coloring matter forms only part of the inert and valueless substances of the crude drugs.

The commercial and officinal fluid preparations of Calisava have been heretofore of the reddish-brown color of the bark. or of an artificial red color-obtained by the addition of cochineal, cudbear, etc.

One of the greatest advances made in pharmacy and medicine has been the separating of the real medicinal qualities of drugs from their inert or repulsive concomitants. Of this the CALISAYA LA RILLA affords a happy example, containing only the pure and total medicinal principles in an absolutely colorless, bright and exquisite cordial.

Calisaya La Rilla is supplied by all druggists.

SOLE MANUFACTURER.

CHAS. ALLEN REED.

9 Cliff Street, New York.

IATROL

IS AN

INODOROUS

Antiseptic Powder

SUPERIOR TO IODOFORM.

A postal card, naming this journal, with your address will bring a large sample free.

Clinton Pharmaceutical Co.

SYRACUSE, N. Y.

mmmm

Manufacturing Chemists.

ESTABLISHED 1866

LENTZ & SONS

SURGICAL INSTRUMENTS.

18 North Eleventh Street,

PHILADELPHIA.

Free! Free!

Our revised and sumptuous catalogue of 500 pages, profusely illustrating all the most desirable devices used in modern surgery, is now ready. A most excellent guide for every physician.

This catalogue will be \$28.00 to \$100.00 mailed free to any physician who contemplates purchasing.



MICROSCOPES

Urinary Analysis

Estimates given on Physicians' ≠ Students'
Outfits.



DEGENERATION OF THE

ispensed by Druggists

HYTOLINE is now recognized by the Medical Profession as being the only Remedy in the Treatment of Obesity that will absorbe the fatty tissue in a great (Walker) z ii degree without any evil after-effects whatsoever

> A REMEDY PREPARED FROM THE ACTIVE PRINCIPLE OF THE BERRIES OF THE PHYTOLACCA DECANDRA

> > For Literature and Clinical Reports address

ST.LOUIS. MO. U.S.A.

ACTIVE PEPSIN The

Webber-Pepsin, s. & D.

A SOLUBLE.

NON-HYGROSCOPIC.

INODOROUS.

PERMANENT PEPSIN.

Dose: 1/8 to 2 Grains.

The Webber-Pepsin will completely digest 6000 times its weight of coagulated egg-albumen in accordance with the conditions of the U.S. P. test or that of the National Formulary.

It is not a Peptone Pepsin, yet it is perfectly soluble, being free from mucus and inert putrescible matter, and is so palatable that the most fastidious patient will take it readily.

It is entirely free from septic contaminations, or other products of decomposition.

Physicians will appreciate these several advantages as possessed solely by the Webber-Pepsin.

Although further investigations based upon the Webber process have enabled us to produce Pepsin capable of digesting from 20,000 to 30,000 times its weight of coagulated egg albumen, we have adopted 1 to 6000 as our standard for the Webber-Pepsin because this already far exceeds the digestive power of any other Pepsin yet introduced, and especially because the production of a Pepsin of a still higher digestive power disproportion-

ately increases its cost.

The Webber-Pepsin is offered in Granular form and in Scales, but unless otherwise specified, we invariably

furnish the GRANULAR form.

Samples furnished upon application.

BRANCH HOUSES: NEW YORK - CHICAGO. Manufactured only by SHARP & DOHME. BALTIMORE, MD., U.S.A.

TRADE

To THE MEDICAL PROFESSION:

An erroneous impression has gone abroad that Peroxide of Hydrogen lacks stability and must of necessity be strongly acid. The reason for this is, that incident to its manufacture many impurities are incorporated in the solution and chemists who are unable to remove these contaminations have created the false idea that their presence prevents loss of oxygen and thereby increases its stability.

The facts are, impurities in Peroxide of Hydrogen render the product unstable by uniting with the oxygen of the solution, and new compounds thus formed are not only irritants and often injurious, but are the cause of the varied and unsatisfactory results so

often experienced by physicians.

A finished product of Peroxide of Hydrogen is not only stable and absolutely harmless, but clinical results obtained from such a product are just as fixed and certain as those from Quinine or Mercury. All chemical and clinical tests prove the O. C. Peroxide of Hydrogen to be a finished product, full 15 vol. strong, free from impurities, non-acid and stable.

Upon receipt of price, 75c. per lb., we will deliver expressage prepaid all orders for 2 lbs. or more.

The Oakland Chemical Co., 89-91 S. Fifth Ave., N. Y.

We will send samples of O. C. Peroxide of Hydrogen and monograph containing instructions for making solutions of H₂ O₂ and glycerine for immediate use, to any physician willing to pay express charges, Physicians should bear in mind that combinations of H₂ O₂ and glycerine after standing one week are of less therupeutic value than pure glycerine.



Glycerine, Sherry Wine, Gentian, Taraxacum, Phosphoric Acid Dil, Carminatives.

The Original Formula of the late John P. Gray, M. D., and used by him for twenty-five years at State Hospital, Utica, N. Y.

An Unequalled Substitute for Cod Liver Oil. Weight and strength often gained faster on the Tonic than on the Oil. Combines the fat-producing elements with a Tonic and Stimulant, and surpasses Malt preparations. Especially useful in Throat troubles of Singers and Speakers. A Superior General and Nerve Tonie. Beneficial effects promptly follow its use in Nervous Prostration, Anæmia, Melancholia, and in all cases of debility following acute exhausting ailments.


An Elegant Pharmaceutical Preparation. Palatable to the taste, creates an immediate appetite, agrees with the most delicate stomach, does not constipate, but stimulates the bowels to healthy action. Invaluable for women and children.

During the Summer Months it is effective in preventing the digestive disturbances so prevalent at that season.

Introduced for the Medical Profession and is endersed by Henry O. Marcy, M. D., Boston; Henry E. Dwight, M. D., Wm. H. Pancoast, M. D., .; J. B. Andrews, M. D., Binghamton, N. Y.; Frank E. Miller, M. D., Willard, N. Y.; Chas. C. Wagner, M. D., Binghamton, N. Y.; Frank E. Miller, M. D., C. M. well, M. D., New York; Wm. T. Cathell, M. D., Geo. J. Preston, M. D., Baltimere; Edwin Everett Smith, M. D., South Wilton, Conn., and others.

Samples, formula and information furnished. Full pint bottle to Physicians, express prepaid, on receipt of one dollar.

Prepared only by THE PURDUE FREDERICK CO., No. 9 Dey Street, near Broadway, N.Y.



\$3.50 Less than Regular Price.

This Regulation 16-inch Black Leather Obstetrical Bag and one year's subscription to FOOD, the new journal of hygiene and nutrition (edited by FRANK WOODBURY. A.M., M D), sent to any address for only

This offer is for a limited time only.

······

Address, THE CLOVER PUBLISHING CO. 71-73 Park Place. New York

Powerful Uterine Sedative.

Particularly efficacious in Engorgement, Inflammation and Induration of the Uterus, in Dysmenorrhoea, Menorrhoea, Leucorrhoea, Amenorrhoea, Prolapsus Uteri, Hysteria, Melancholia, Vomiting of Pregnancy, Habitual Abortion, and Uræmic Eclampsia.


It is not a nauseous compound, but an elegant elixir. It is a practical Emmenagogue, a safe and reliable Parturient, Antispasmodic, Diuretic, and Tonic.

FORMULA -Caulocorea is composed of the active principles of Caulophyllum, Thalictroides, Viburnum Opulus, Prunifolium, Dioscorea Villosa, Mitchella Repens, Aletris Farinosa, combined with Spts. Ætheris Comp. and Aromatics.

A Pound Bottle—Free

to any physician who will pay express charges and say this journal induced him to write. It is prepared solely for the use of physicians, and advertised only in medical journals.

CAULOCOREA MANUFACTURING CO., Portland, Me.



HIGH OR LOW TEMPERATURE?

Very few physicians have access to Milk Laboratories, such as described by Dr. T. M. Rotch, of Boston, in Archives of Pediatrics, Feb., 1893, or by Dr. Henry Koplik in the New York Medical Journal, Feb. 4th, 1893, but just as reliable results can be obtained in every family by the use of an

ARNOLD STEAM STERILIZER.

The Cover of this apparatus is now so constructed that a temperature of 100°C. is maintained with the Hood on, and δ_5 °C. to 90°C. with it off.

This simplifies and solves the problem of infant feeding, under the physician's direction, as far

as scientific knowledge has reached.

Always prescribe "Arnold Sterilizer with Perforated Cover." It meets every requirement. For sale by Druggists and Instrument Dealers.

Write direct to the manufacturers for full information.

WILMOT CASTLE & CO., Rochester, N. Y.

We also manufacture Sterilizers for all other purposes—steam, boiling water, and hot air.

ECZENIA

IT IS SAID THAT 70 PER CENT. RECOVER WITH "VINOLIA," AND ALL ARE RELIEVED.

"Vinolia" Cream is cooling and soothing for eczematous and other irritable and irritated conditions of the skin as well as mucous membranes. It is a plastic, homogeneous antisep-tic, containing "VINOLIA" BASIS (a milled basis composed of simple and intimately associated pure fatty bodies) and a very small proportion of a levigated form of zinc, camphor, and scents, but no cocaine, morphine, belladonna, etc. It is absolutely free from fatty acids and from all objections to be brought against ointments made with mineral and other bases. It is bland, innocuous, emollient, and agrees in every way with the most delicate and diseased skins. It is cleanly to use, and all who prescribe it pronounce it a pleasant and extremely agreeable "cream"—indeed, an exquisite toilet

preparation as well as a grateful application.
"Vinolia" will relieve the itching of eczema, of insect bites, chilblains, and various skin affections as soon as applied. Cases of eczema lasting many years have proved amenable to treatment with *Vinolia*, being cured by it in a short time. We have in our possession a very large number of letters from medical men (some of them leading dermatologists) attesting in no uncertain terms to the unquestionable usefulness of VINOLIA for itching and skin irritation. Very many report favorably after numerous trials made with it in acne rosacea. It has been used with noteworthy advantage for lying-in women, sore nipples, gynæcological examinations and treatment, for urticaria, cold sores, chapped hands, redness, freckles, and sunburn, no matter how produced. Its grateful action renders it unsurpassable in piles and pruritis ani, herpes, styes, blepharitis, intertrigo, irritable abraded ulcers, and prickly heat. Its application to the face in cases of prospective exposure or after shaving is a salutary measure for delicate skins. It agrees with children admirably. "VINO-LIA" CREAM is perfectly preservable, harmless to the skin, and renders this tissue elastic, smooth, supple, and natural. It is of service in facial eruptions, which are so often caused and kept up by deleterious cosmetics and hardening soaps. It readily penetrates the outer layer of the skin; is perfectly innocuous, anti-inflammatory, and applicable in all states of the integument. Where there is thickening of the epidermis, and where there are crusts, VINOLIA fulfils all requirements. It has the agreeable odor of the French rose, and forms a striking contrast to the malodorous, rancid ointments so often seen. We repeat, it will relieve the most intractable itching as soon as applied.

DIRECTIONS FOR USE.

VINOLIA may be very freely used at any and all times. Irritable surfaces and delicate skins may be washed with warm water and "VINOLIA" SOAP, dried gently with a soft linen towel or silk handkerchief, and covered with a thin layer of "VINOLIA," which should be renewed, where practicable, at least thrice daily

REPORTS ON VINOLIA. SIX CASES OF ECZEMA.

GEO. P-, Esq., M. D., reports :- "I have prescribed Vinolia in about half-a-dozen cases of eczema, in varying rior to all others I have before recommended."

situation and severity, and in all of them have had reason to be satisfied with the results."

ECZEMA AND ACNE.

M. D. reports:—"I have used Vinolia always with good results, especially in cases of eczema and acne."

M. D. reports:-" I have made extensive use of Vinolia, and find it more valuable every day. My latest use of it has been to allay the irritation of a part which had been painted with tincture of iodine, which it did most effectually. In a case of inflamed acne rosacea, I have also proved it a great comfort. I fancy that its range of usefulness will increase as it becomes better known.

E. B. F—, Esq., M. D., reports:—"I have tried Vinolia in cases of acne of the face, and in a case of eczema solare I found it a very efficacious remedy."

THE NEW SUPER-FATTED SOAP.

Medical Men, and those with skin diseases and delicate skin, know only too well the want of a soap for medical and toilet purposes free from alkaline properties. "VINOLIA" SOAP supplies the requirement, it is a perfect super-fatted soap. It differs wholly from other soaps, which, by reason of their alkaline nature, are ofttimes harmful in skin affections, inasmuch as it contains, instead of some free alkali, an excess of plastic emollient cream. It is unadulterated, and suits the very tenderest skin. As a detergent it cannot be surpassed, and it is tonic, antiseptic, emollient, and pleasant. "VINO-LIA" SOAP does not dry the skin like other soaps, as the presence of the extra cream prevents the possibility of any free alkali, even in the water, attacking the integument, extracting the oil, and so hardening and weakening the tissue, thus rendering it liable to eruptions and premature decay. The deleterious action of ordinary soaps upon the hair and scalp is easily noticeable after shampooing with them, and their retarding effect in treatment is known to all dermatologists. "VINOLIA" SOAP lathers beautifully, and the difference between its effect on the skin and that of other soaps will appear the moment a tablet is used. It is the only soap for skin treatment, and for the toilet, nursery, and bath. In ecsema it is the greatest possible desideratum. Those who have used it once use no other.

REPORTS ON VINOLIA SOAP.

W. W—, Esq., M. D., reports:—"There can be no doubt about Vinolia being a most charming preparation, and the 'Vinolia' Soap is without exception the most agreeable I have ever used."

T. R. A .---, Esq., M. D., reports:--"I find 'Vinolia' Soap the best soap I know of in all skin eruptions, or in cases where the skin is tender, as it does not irritate. I find it supe-

PRICES:

- "VINOLIA" CREAM (for Itching, Face Spots, Eczema, etc.), 50 cts., \$1.00, and \$1.75.
- "VINOLIA" POWDER (for the Toilet, Nursery, Roughness, etc.), 50 cts., \$1.00, and \$1.75.
- "VINOLIA" SOAP, Floral, 15 cts.; Medical, 25 cts.; Toilet, 35 cts.; and Vestal, 85 cts.

Samples and Pamphlets giving full Medical and Analytical Reports free on application. "Vinolia" Soaps and Preparations may be obtained of any good druggist.

BLONDEAU et CIE., 73-77 Watts Street, New York.

SYR. HYPOPHOS. CO., FELLOWS

Contains the Essential Elements of the Animal Organization—Potash and Lime;

The Oxidising Agents—Iron and Manganese;

The Tonics—Quinine and Strychnine;

- And the Vitalizing Constituent—Phosphorus; the whole combined in the form of a Syrup with a Slightly Alkaline Reaction.
- It Differs in its Effects from all Analogous Preparations; and it possesses the important properties of being pleasant to the taste, easily borne by the stomach, and harmless under prolonged use.
- It has Gained a Wide Reputation, particularly in the treatment of Pulmonary Tuberculosis, Chronic Bronchitis, and other affections of the respiratory organs. It has also been employed with much success in various nervous and debilitating diseases.
- Its Curative Power is largely attributable to its stimulant, tonic and nutritive properties, by means of which the energy of the system is recruited.
- Its Action is Prompt; it stimulates the appetite and the digestion, it promotes assimilation, and it enters directly into the circulation with the food products.
- The prescribed dose produces a feeling of buoyancy, and removes depression and melancholy: hence the preparation is of great value in the treatment of mental and nervous affections. From the fact, also, that it exerts a double tonic influence, and induces a healthy flow of the secretions, its use is indicated in a wide range of diseases.

NOTICE-CAUTION.

The success of Fellows' Syrup of Hypophosphites has tempted certain persons to offer Imitations of it for sale. Mr. Fellows, who has examined samples of several of these, finds that no two of them are identical, and that all of them differ from the original in composition, in freedom from acid reaction, in susceptibility to the effects of oxygen when exposed to light or heat, in the property of retaining the strychnine in solution, and in the medicinal effects.

As these cheap and inefficient substitutes are frequently dispensed instead of the genuine preparation, physicians are earnestly requested, when prescribing the Syrup, to write, "Syr. Hypophos. Fellows."

As a further precaution, it is advisable that the Syrup should be ordered in the original bottles; the distinguishing marks which the bottles (and the wrappers surrounding them) bear, can then be examined, and the genuineness—or otherwise—of the contents thereby proved.

Medical Letters may be addressed to:



TRADE MARK.

PREVENTS FERMENTATION OF FOOD IN THE STOMACH. MOST POWERFUL REMEDY FOR HEALING PURPOSES. CURES:

Send for free book of 80 pages giving articles by contributors to medical literature. PHYSICIANS WILLING TO PAY EXPRESS CHARGES WILL RECEIVE FREE SAMPLES ON APPLICATION.

Glycozone is sold only in 4 ounce, 8 ounce, and 16 ounce bottles. Never sold in bulk.

CH. MARCHAND'S

PEROXIDE OF HYDROGEN. (MEDICINAL) Ha Os

ENDORSED BY THE MEDICAL PROFESSION. USED BY THE HOSPITALS OF THE U.S. ARMY.

SOLD BY LEADING DRUGGISTS.

Prepared only by

Chemist and Graduate of the "Ecole Centrale des Aris et Ma (France)

Mention this publication.

Laboratory, 28 Prince St., New York.



COLDEN'S LIEBIG'S LIQUID EXTRACT OF BEEF AND TONIC INVIGORATOR. LABEL.

ESSENTIALLY DIFFERENT FROM ALL OTHER REEF TONICS. UNIVERSALLY ENDORSED BY LEADING PHYSICIANS.

This preparation, consisting of the Extract of Beef (prepared by Baron Liebig's process), the best Brandy obtainable, soluble Citrate of Iron, Cinchona and Gentian is offered to the Medical Profession upon its own merits. It is of inestimable value in the treatment of Debility, Convalescence from Severe Illness Ansemis, Malarial Fever, Chilorosis, Incipient Consumption, Nerveus Weakness, and maladies requiring a Tonic and Nutrient. It is quickly absorbed by the Stomach and upper portion of the Alimentary Canal, and therefore finds its way into the circulation quite rapidly.

COLDEN'S LIQUID BEEF TONIC appeals to the judgment of intelligent Physicians in the treatment of

ALL CASES OF GENERAL DEBILITY.

By the urgent request of several eminent members of the medical profession, I have added to each wineglassful of this preparation two grains of Soluble Citrate of Iron, and which is designated on the label, "With Iron, No. 1;" while the same preparation, Without Iron, is designated on the label as "No. 2." In prescribing this preparation, physicians should be particular to mention "COLDEN'S," viz., "Est. Carate Ft. Comp. (Colden.")" A Sample of COLDEN'S BERF TONIC will be sent free on application, to any physician (enclosing business card) in the United States. Sold by druggists generally.

C. N. CRITTENTON, General Agent, 115 Fulton St., New York.

Constantine's Pine-Tar Soap.

Wholesale Depot, C. N. CRITTENTON, 115 Fulton St, New York. Samples of above Soaps SENT FREE, on application, to any Physician enclosing card.

FOR THE PREVENTION AND CURE OF

PULMON PHTHISIS.

FORMULÆ

COUGH TABLETS.

EACH TABLET CONTAINS

Morph. Sulph. (18 gr.), Atropiæ Sulph. (18 gr.), Codeia (18 gr.), Antimony Tart. (18 gr.), Ipecac, Aconite, Pulsatilla, Dulcamara, Causticum, Graphite, Rhus-tox, and Lachesis, fractionally so arranged as to accomplish every indication in any form of cough.

CONSTITUENT TABLETS.

EACH TABLET CONTAINS

Arsenicum (10 gr.), Precipitate Carb. of Iron, Phos. Lime, Carb. Lime, Silica, and the other ultimate constituents, according to physiological chemistry (normally), in the human organism, together with Caraccas, Cocoa and Sugar.

PRICE, THREE DOLLARS PER DOUBLE BOX.

Containing sufficient Tablets of each kind to last from one to three months according to the condition of the patient.

SPECIAL OFFER.

While the above formulæ have been in use, in private practice, over 30 years, and we could give testimonials from well-known clergymen, lawyers and business men, we prefer to leave them to the unbiased judgment of the profession with the following offer: On receipt of 50 cents, and card, letter-head, billhead, or other proof that the applicant is a physician in active practice, we will send, delivered, charges prepaid, one of the regular (double) boxes (retail price, Three Dollars), containing sufficient of each kind of Tablets to test them three months (in the majority of cases), in some one case. Card, letter-head, or some proof that the applicant is a physician in active practice, MUST accompany each application. Pamphlet, with full particulars, price list, etc., on request.

As we furnish no samples through the trade, wholesale or retail, for samples, directions, price lists, etc., address

I. O. WOODRUFF & CO.,

Manufacturers of Physicians' Specialties.

38 MAIDEN LANE.

NEW YORK



At the National Capital,

which is another way of saying, "in our own Country," where prophets usually have no honor,

All Physicians Prescribe

As the most invaluable agent in the treatment of Coughs, Catarrh, Phthisis, Pulmonalis, Tuberculosis, General Debility, etc.

A purified and tasteless preparation of Petroleum for internal use—a substitute for Cod Liver Oil and Emulsions. A distinct point in its favor is that it will not cause nausea or eructations, and the most delicate stomach can tolerate it and grow strong in its use. It is not a patent medicine in any sense of the word, but is sold on the prescription of the practitioner, and it does not end in Promising to Cure, but does its whole duty to the entire satisfaction of the attending physician and ailing patient. It is worth a trial

For Coughs, Consumption, etc.


and for such we will send a dollar bottle free to any physician who will pay express charges.

Manufactured only by

1316 L Street, N. W., Washington, D. C.

SOLD BY ALL DRUGGISTS

ONE DOLLAR A BOTTLE



Chnical evidence shows that apart from the value of Cod Liver

Dr. CHAS. W. MING. Medical Sant

Oil as an assimilable fat, the active principles of the oil, known collectively as Morrhuol, represent its true therapeutic value.

Morrhuol obtains excellent results wherever the oil is indi-

cated, but is not

ORRHUOL

(CHAPOTEAUT.)

The Alkaloids and all the active principles of Cod Liver Oil dispensed in spherical capsules, each of which is equivalent to one teaspoonful of oil.

tolerated. It is the type of those remedies used in wasting diseases which act by stimulating nutrition.

Morrhuol Creosote is recom-

mended for the treatment of bronchial catarrh. and tuberculosis. The stomach rapidly accomodates

MORRHUOL

In spherical capsules, each of which contains I minim of Creosote (equal to 90% of Guaiacol) and 3 minims of Morrhuol.

Dose: 4 to 12 capsules daily.

crease in the appetite and improved general condition, besides decrease in the Expectoration,

a noticeable in-

Night sweats and Cough.

After a few days there is

itself to large doses.

RIGAUD & CHAPOTEAUT, Paris,

New York Depot: E. FOUGERA & CO., 30 N. William Street.

From whom samples and literature may be obtained on application

DR. WILLIAM A. HAMMOND'S PRIVATE HOSPITAL

For DISEASES of the NERVOUS SYSTEM.

WASHINGTON, D. C.

The Hospital is situated on Columbia Heights, at the corner of Fourteenth Street and Sheridan Avenue. The position is the highest in the immediate vicinity of Washington. The soil is dry and all the surroundings are free from noxious influences. The Fourteenth-street cable cars run almost to the door. The building is very large, and in its construction every sanitary requirement has been fulfilled. The number of patients is limited to twenty. Electricity in all its forms, baths, douches, massage, inhalations, nursing, etc., are provided as may be required by patients, in addition to such other medical treatment as may be deemed advisable. The building is heated throughout by steam. For further information Dr. Hammond can be addressed at the



HOSPITAL, 14th STREET and SHERIDAN AVENUE, WASHINGTON. D. C.

PRIVATE HOSPITAL

BEECHHAVEN,

NEW PARIS, OHIO.

For Nervous and Mental Diseases, Opium, Morphine and Chloral Habits.

Under the personal supervision of Dr. Charles W. King, for six years past Superintendent of the Dayton Asylum for the Insane.

New buildings, most modern improvements. Sixty acres of lawn. The "Castalia" Mineral Spring immediately on our grounds. Address

Dr. CHAS. W. KING, Medical Supt.,

(On P., C. & St. L. R. R.)

NEW PARIS, PREBLE CO., OHIO.

BURN BRA



A PRIVATE HOSPITAL

FOR MENTAL AND NERVOUS DISEASES.

Founded by the late Robert A. Given, M. D., in 1850.

Extensive and beautiful grounds. Perfect privacy. A pleasant, safe and healthful home. Music, games, open-air amusements. The oldest institution of the kind in the United States. Both sexes received.

ARRANGEMENTS MADE FOR CHRONIC CASES.

Located a few miles west of Philadelphia, at Primos Station, on the P. W. REFERENCES:

Professors H. C. Wood, Wm. Pepper, Alfred Stillé, William Goodell, Roberts Bartholow, R. A. F. Penrose, J. M. DaCosta, Charles K. Mills, James Tyson, and Dr. Lawrence Turnbull: Prof. William Osler, of Johns Hopkins University; W. W. Lassiter, M. D., Petersburg, Va. RESIDENT PHYSICIANS:

J. Willoughby Phillips, M. D., For further information address S. A. Mercer Given, M. D.

BURN BRAE, Clifton Heights, Delaware Co., Pa.

PURIFIED OPIUM.

47-FOR PHYSICIANS' USE ONLY. TA

Contains the Anodyne and Soporific Alkaloids, Codeia, Nar-ceia and Morphia.

Excludes the Poisonous and Convulsive Alkaloids, Thebaine, Narcotine and Papaverine.

SVAPNIA has been in steadily increasing use for over twenty years, and whenever used has given great satisfaction.

To Physicians of Repute, not already acquainted with its merits, samples

will be mailed on application.

Svarnia is made to conform to a uniform standard of Opium of Ten per cent. Morphia strength.

JOHN FARR, Manufacturing Chemist, New York.

C. N. CRITTENTON, Gen'l Agent, 115 Fulton St., N. Y. To whom all orders for samples must be addressed.

SVAPNIA IS FOR SALE BY DRUGGISTS GENERALLY.

DR. MARTIN'S VACCINE VIRUS.

PRICE REDUCED.

PHICE REDUCED.

10 Large Ivory Points, \$1.00. Fully warranted.
Boards of Health, and on large orders. In 1870 we introduced practice of animal vaceination. Our establishment, continued since, is by far the Oldest, Largest and Best Appointed in the cou hitherto the most expensive, can now be obtained by the profeprice as any other. No virus ours unless package bears our signature.

our signature. **
DR. HENRY A. MARTIN & SON

Brookline Station, Boston, Mass.

UNIVERSITY OF PENNSYLVANIA, MEDICAL DEPARTMENT.

The 188th Annual Winter Session will begin Monday, October 2, 1893, at 12 M., and will end at Commencement, June 5, 1894.

The PRELIMINARY and SPRING SESSIONS have been discontinued.
The Curriculum is graded, and attendance upon four annual Winter Sessions is required. College graduate in Arts or Science, who have pursued certain Biological studies, are admitted to advanced standing.
Practical Instruction, including laboratory work in Chemistry, Histology, Osteology, and Pathology, with Bedside Instruction in Medicine, Surgery, Gynaecology and Obstetrics are a part of the regular course, and without additional expense.
For Catalogue and announcement containing particulars, apply to

36th St. and Woodland Ave., Phila. DR. JOHN MARSHALL, Dean.



The Interesting Offer

Of a handsome double morocco pocket medicine case—7 1/8 x 3 3/4 x 7/8 inches—containing 24 vials filled with a complete assortment of our popular Tablets and Triturates at the special price of \$3.00 has been

The case contains:

Acetanalide Comp. (Migraine).
Anti-Constipation (Mulford Co.)
Chlorodyne (Mulford Co.)
Viburnum Comp. (valuable uterine tonic).
Ammonii Chloridum Comp.
Bismuth et Cereii Ozalat Ammonii Chloridum comp.

Bismuth et Cereii Ozalat
Bronchitis
Cathartic, "Active,"
Diarrhoea.
Ergotine (Bonjean's), z grain,
Ergotine (Bonjean's), z grain,
Fevre t Acid, Arsen. et Strych, No. z.
Fever (Dr. T. G. Davis),
Hydrarg, Chlor. Mite et Ipecac et Sodii Bicarb, No. z.
Hydrarg, Iod. Viride, z-4 grain,
Morphina Sulphas, z-6 grain,
Neuralgic (Dr. H. J. Kenyon),
Nitro-Glycerin Comp. (Mulford Co.)
Pulv. Ipecac et Opii, (U. S. P.) representing 2½ grains,
Tinct. Acontit, z minin., z grain (colored blue, antiseptic).
Quinines Sulphat, z grain.
Hydrarg. Chlor. Mite, z-10 grain,
Kermes Mineral Comp. (Dr. E. Wilson).

Accepted by over 15,000 Physicians

These cases are a permanent advertisement for us. as they are constantly being refilled with our standard preparations, which physicians use because they are so reliable, valuable, convenient and elegant-all compressed. We continue this attractive \$3.00 offer for a limited period. Send check, money-order or postal note. We deliver it free of postage.

H. K. MULFORD COMPANY, 2132 Market Street, Philadelphia



The Demand For

a pleasant and effective liquid laxative has long existed—a laxative that would be entirely safe for physicians to prescribe for patients of all ages—even the very young, the very old, the pregnant woman and the invalid—such a laxative as the physician could sanction for family use because its constituents were known to the profession and the remedy itself had been proven to be prompt and reliable in its action, as well as pleasant to administer and never followed by the slightest debilitation. After a careful study of the means to be employed to produce such ployed to produce such

A Perfect Laxative

the California Fig Syrup Company manufactured, from the juice of True Alexandria Senna and an excellent combination of carminative aromatics with pure white sugar, the laxative which is now so well and favorably known under the trade name of "Syrup of Figs." With the exceptional facilities, resulting from long experience and entire devotion to the one purpose of making our product unequaled, this demand for the perfect laxative

is met by Our Method

of extracting the laxative properties of Senna without retaining the griping principle found in all other preparations or combinations of this drug. This method is known only to us, and all efforts to produce cheap imitations or substitutes may result in injury to a physician's reputation, and will give dissatisfaction to the patient; hence, we trust that when physicians recommend or prescribe "Syrup of Figs" (Syr. Fici Cal.) they will not permit any substitution. The name "Syrup of Figs" was given to this laxative, not because in the process

of Manufacturing

a few figs are used, but to distinguish it from all other laxatives, and the United States Courts have decided that we have the exclusive right to apply this name to a laxative medicine. The dose of

"SYRUP OF FIGS"

as a laxative is one or two teaspoonfuls given preferably before breakfast or at bed time. From one-half to one tablespoonful acts as a

as a laxative is one or two teaspooning given precising years and seed at the purgative, and may be repeated in six hours if necessary.

"Syrup of Figs" is never sold in bulk. It is put up in two sizes to retail at fifty cents and \$1.00 per bottle, and the name "Syrup of Figs" as well as the name of the California Fig Syrup Company is printed on the wrappers and labels of every bottle.

CALIFORNIA FIG SYRUP COMPANY.

SAN FRANCISCO, CAL.

LOUISVILLE, KY.

NEW YORK, N. Y.

THE POST-GRADUATE

The June Number of this Monthly has just been issued. The following is the TABLE OF CONTENTS:

THE MONTH.

Portrait of the late Professor CHARLES CARROLL LEE, M. D., LL. D.

The late Professor Charles Carroll Lee-His Place as a Teacher made in the Post-Graduate-The Unclean Condition of the Land Bordering on our Railway Thoroughfares-Infection of the Water Supply through want of Cleanliness of the Soil-Political Street Cleaners-The Secret of Keeping the Streets Clean is not to Let the Dirt Accumulate-New York Much Cleaner than Chicago-A General Law Exempting Educational and Medical Institutions from Taxation-Electricity as a Disinfectant-Sir Thomas Brown and Incurable Disease-Epileptics and their Strong Appeal to Human Sympathy-The Deplorable Consequences of Epilepsy-Colonization and Segregation of Epileptics-The Governor has Vetoed the Bill for a Colony in this State-His Reasons for this Action-A Paradise-deserving Philanthropy-The Progress of the New Building.

LEADING ARTICLES.

Notes of a Case of Cancer of the Uterus: Remarks on Diagnosis and Treatment, by Daniel Lewis, M. D.-The Therapeutic Value of the Oily Sprays and the various Powders now used in Nasal and Throat Diseases, as Compared with Older Remedies, by Clarence C. Rice, M. D.

VARIA.

The New York Academy of Medicine, April 20th, 1893. Discussion of Dr. Lewis' Paper by Dr. Hanks-Extract of Letter from Dr. George E. Abbott.

THE CLINICAL SOCIETY.

Report of a Case, by Robert P. Oppenhimer, M. D., and Discussion-Dr. Brannan, Dr. Leszynsky, Dr. Porter-Discussion of Dr. Davis' Paper on Chronic Suppurative Otitis Media of Tubercular Origin: Dr. Brannan, Dr. Frank N. Lewis, Dr. Davis-Presentation of Specimens by Dr. J. W. Brannan and Discussion: Dr. Einhorn, Dr. Lewis.

It is an Independent Journal, containing only Original Articles and Notes on Current Medical Topics of interest to the whole Medical Profession in this Country.

SUBSCRIPTIONS, \$1.00 A YEAR.

SINGLE NUMBERS, 20 CENTS.

Address "THE POST-GRADUATE," No. 226 East 20th Street, New York City.

QUIZZING.

Physicians prepared for Army, Navy, Marine-Hospital and General Hospital EXAMINATIONS:

FOR THE REGENTS' EXAMINATION PRIOR TO BEGINNING PRAC-TICE IN THE STATE OF NEW YORK.

IN ADDITION.

SPECIAL PRACTICAL COURSES

Including Operative Surgery on the Cadaver, Minor Surgery and Bandaging, and Physical Diagnosis.

These practical courses may be taken separately by other physicians than our students. Address

Drs. Dawbarn & Symonds

105 West 74th Street, NEW YORK.

NEW YORK POLYCLINIC AND HOSPITAL.

A Clinical School for Practitioners of Medicine. Physicians may enter at any time. Sessions continuous throughout the year. For catalogue or information address

L. EMMETT HOLT, M. D., Secretary, 214-18 East Thirty-fourth Street, New York.

PHILADELPHIA POLYCLINIC.

Post-Graduate Teaching in six-weeks' and three-months' courses actual Clinical Work, with abundant material and small classes.

DR. A. W. WATSON,

18th and Lombard Streets, PHILADELPHIA.

CHICAGO POLICLINIC ≠ HOSPITAL 174-176 CHICAGO AVENUE.

A CLINICAL SCHOOL FOR PRACTITIONERS OF MEDICINE

Physicians can enter at any date. Entire Faculty on duty throughout the summer. The Building is the largest and most thoroughly equipped for Post-Graduate instruction in the West. The Hospital contains soo beds. The Clinical material is abundant. Situated in the same section of the city are several large hospitals to which our matriculates only are admitted. Specia: clinics, lectures and demonstrations will be given by distinguished medical men from abroad during the World's Fair season.

For information address the Corresponding Secretary,

MORE 11 P. BROWN, 174-175 Chicago Ave., Chicago, Ill.

POST-GRADUATE MEDICAL SCHOOL

OF CHICAGO.

819, 821, 823 West Harrison St., Chicago, Ill.

A school for practitioners of medicine. Tickets may be taken out with equal advantage at any time. Write for Bulletin No. 4, for Summer Course and Special World's Fair Course of Lectures, to

FRANKLIN H. MARTIN, Sec'y.

THE COLLEGE OF PHYSICIANS AND SURGEONS BALTIMORE, MD.

Offers unexcelled facilities for acquiring a thorough Scientific and Practical Medical Education.

NEW COLLEGE BUILDINGS, WELL-EQUIPPED LABORATORIES, THREE LARGE GENERAL HOSPITALS, LYING-IN ASYLUM.

Next Regular Session begins October 2, 1893. For Caralogue giving detailed information address

PROF. THOS. OPIE. M.D., DEAN, Baltimore, Md. 600 N. Howard Street.

BELLEVUE HOSPITAL MEDICAL COLLEGE.

CITY OF NEW YORK.

SESSIONS OF 1893-94.

The REGULAR SESSION begins on Monday, September 25, 1893, and continues for twenty-six weeks. During this session, in addition to the regular didactic lectures, two or three hours are daily allotted to clinical instruction. Attendance upon three regular courses of lectures is required for graduation. The examinations of other accredited Medical Colleges in the elementary branches are accepted by this College

The Spring Session consists of daily recitations, clinical lectures and exercises and didactic lectures on special subjects. This session begins March 26, 1894, and continues until the middle of June.

The CARNEGIE LABORATORY is open during the collegiate year, for instruction in microscopical examinations of urine, practical demonstrations in medical and surgical pathology, and lessons in normal histology and in pathology, including

For the annual Circular, giving requirements for graduation and other information, address

tion, address
PROF. AUSTIN FLINT, Secretary,
Bellevue Hospital Medical College, foot of E. 26th St., NEW YORK CITY.

UNIVERSITY OF MARYLAN

SCHOOL OF MEDICINE.

Session of 1893-94.

PACULTY:

GEO. W. MILTENBERGER, M. D.,

SAMUEL C. CHEW, M. D.,
Professor of Principles and Practice of Medicine
and Clinical Medicine.

WILLIAM T. HOWARD, M. D., Prof. of Diseases of Women and Children, and Clinical Medicine.

JULIAN J. CHISOLM, M. D., or of Eye and Ear Di

FRANCIS T. MILES, M. D., esor of Physiology, and Clinical Pro of Diseases of the Nervous System.

J. EDWIN MICHAEL, M. D., Professor of Obstetz

I. EDMONDSON ATKINSON, M.D., Professor of Materia Medica and Therapeutics Clinical Medicine and Dermatology.

R. DORSEY COALE, Ph. D., RANDOLPH WINSLOW, M. D.,

JNO. NOLAND MACKENZIE, M.D., L. McLANE TIFFANY, M. D., Clinical Professor of Diseases of the Threat and Noss.

Next regular session begins October 2, 1893. For catalogue and other information, address

J. EDWIN MICHAEL, M. D., Dean, 201 W. Franklin St., Baltimore, Md.

DENTAL DEPARTMENT.

FACULTY:

F. J. S. GORGAS, M. D., D. D. S.,

JAS. H. HARRIS, M. D., D. D. S.,

or of Operative and Clin I. E. ATKINSON, M. D.,

of Materia Medica a FRANCIS T. MILES, M. D., Professor of Physiology

L. McLANE TIFFANY, M. D., Ctinical Profes nor of Oral Surgery.

RANDOLPH WINSLOW, M. D., Professor of Angle

R. DORSEY COALE, Ph. D., Professor of Chemistry.

Session opens October s, 1893. For catalogue or other information, address

Dr. FRED. J. S. GORGAS, Dean, 845 Eutaw St., Baltimore, Md.

SOCIETY OF THE LYING-IN HOSPITAL OF THE CITY OF NEW YORK.

THE MIDWIFERY DISPENSARY has been absorbed by this institution. The number of deliveries during 1892, 9,55; during 1892, 2,070. The administrative building is in the most densely populated tenement-house district in the city. There are accommodations for twenty students, the regular term of instruction being two weeks, with six to fifteen deliveries for each students. Instruction is given during the entire year and is open to practitioners of medicine and students who have attended one course of medical lectures. Students are admitted in the order in which their applications are received, or they may be assigned to special dates when attended one course or mecutar fectures. Students are admitted in the order in which their applications are received, or they may be assigned to special cases when practicable. While waiting they can attend the various hospital clinics, dispensaries, etc., in the city. All deliveries are strictly antiseptic, and all cases are attended at their houses as in private practice. Lodgings, instruments and medicines are furnished at the hospital during the two weeks' service, and certificates are issued to those who have performed the service satisfactorily. [Separate apartment for graduates] For further information apply to CHAS. FORD, Superintendent, 314 Broome St., New York City. Special instruction upon the manikin given by the attending physicians.

UNIVERSITY OF MICHIGAN.

DEPARTMENT OF MEDICINE AND SURGERY.

Forty-fourth Year, commencing Monday, Oct. 2, 1893.

The Course of Instruction now comprises four collegiate years of nine monthle each. Examination for admission required. Women admitted.

FEBS:—For the first year—Residents of Michigan, \$35; non-residents, \$60.

For each subsequent year—Residents of Michigan, \$35; non-residents, \$35.

Graduation Fee for all alike, \$10.

Circular and Catalogue, with full details, sent on application.

WM. A. CAMPBELL, M. D., Secretary,

Ann Arbor, Michigan.

BALTIMORE MEDICAL COLLEGE

PRELIMINARY FALL COURSE begins September 1, 1893.
REGULAR WINTER COURSE begins October 1, 1893.
EXCELLENT TEACHING FACILITIES, CAPACIOUS HOSPITAL, LARGE CLINICS.

Send for catalogue and address

DAVID STREETT, M. D., Dean, 403 N. Exeter St., Baltimore, Md.

Northwestern University Medical School. N. S. DAVIS, M. D., LL. D., DEAN.

The course is graded and four annual sessions are required. Special arrangements are made for students who have taken the courses antecedent to the study of medicine in universities or colleges.

Laboratory work in Anatomy, Embryology, Normal and Pathological Histology, and Chemistry, is compulsory and without special charge.

Clinical work is obligatory and the class is divided into a number of small sections. The material is ample.

For a circular of information address the Secretary,

DR. FRANK BILLINGS, 235 STATE STREET, CHICAGO, ILL.

The Baltimore University School of Medicine The Regular Course of Lectures in this Institution will begin October 2, 1893.

Clinical teaching a prominent feature. Clinics are held daily during the ourse and include General Medicine, Dermatology, Eye and Ear, Gynecology, biseases of the Nervous System, Diseases of Children, Surgery, and Diseases of hroat and Chest. For Catalogue and further information, address

E. W. EILAU, M. D., DEAN, 1523 E. Baltimore Street.

WESTERN PENNSYLVANIA MEDICAL COLLEGE.

Medical Department of the WESTERN UNIVERSITY OF PENNSYLVANIA.

RIGHTH ASPLIA. REQUIAN SERVICE Spiles on the last Tureday of September. 1805, and continues dis mostles Astondance upon three require covers of Lectures is requisite for graduation. A three years' graded course is provided. The Spiles Service Segment begins the second Tureday is April, 1894 and continues ten weeks.

Special importance attaches to "the superior clinical advantages possessed by this College." For particulars, see annual announcement and catalogue, for which address the Secretary

Prof. T. M. T. McKENNAN, 810 Penn Avenue.

Business Correspondence should be addressed to
Prof. W. J. ASDALE, 2107 Penn Avenue, Pittsburgh, Pa.

Medical Department of the Columbian University, Washington, D. C. Medical Department of the Columbian University, Washington, D. C.

Announcement 1893-94. The Seventy-second Session will begin Oct. 1, '93, and continue 7 months. Graded 4-year course required. Ample clinical facilities. FACULTY-1. F. Thompson, Surg.; W. W. Johnson, Prac.; A. F. A. King, Obstetrics; F. A. de Schweinitz, Chemistry; Wm. P. Carr, Physiology; D. W. Prentiss, Therapeutics; D. K. Shute, Anatomy; H. C. Yarrow, Dermatology; G. B. Harrison, Pediatrics; Theo. Smith, Bacteriology and Hygiene; T. E. McArdle, Minor Surgery; H. L. E. Johnson, Gynzcology; G. N. Acker, Pathological Histology; W. M. Gray, Normal Histology; W. K. Butler, Ophthalmology; S. Ruffin, Medical Jurisprudence; C. W. Richardson, Laryngology and Otology; A. C. Patterson, Mental Disease; E. L. Tompkins, Neurology.

For circulars, address

A. F. A. King, M. D., Dann,
1315 Massachusetts Avenue, N. W., Washington, D. C.



TRY IT FOR INDIGESTION AND DYSPEPSIA.

The amount of study given to gastric disorders and papers written on the same since the time of Trousseau, by the medical profession, would make a small library; yet, notwithstanding all this research, troublesome cases are daily met with by the regular practitioner, and while the majority of physicians fully understand the physiology of digestion, and the pathology and therapeutics of stomach disorders, and are fully able to review the causes of indigestion, and the symptoms and clinical varieties of Dyspepsia, yet the present mode of treatment is very unsatisfactory. Why not go back to first principles, and try Burnham's Clam Bouillion?

WHAT IS IT?

It is the concentrated juice or extract of the quohaug or clam, put up in bottles and sterilized. It will keep in any climate. It is more strengthening than milk or extract of beef. It is easily prepared by simply adding hot water. You will find it a very acceptable change from the long list of Beef and other food products now offered. Put up in three sizes, 25-cent, 50-cent and \$1.00, and is sold by all first-class Druggists and Grocers. Specify Burnham's Clam Bouillon, which is put up only in glass bottles. A sample bottle will be sent free to any physician upon application to the

E. S. BURNHAM COMPANY,

120 Gansevoort St.,

NEW YORK.

***KUMYSGEN**



Or Kumyss in powder form for making Liquid Kumyss in less than one minute, by simply dissolving the powder in water.

More nutritious and more palatable than any Liquid Kumyss.

There is no Food that equals it in all forms of Indigestion, Pulmonary Affections, Fevers, Vomiting in Pregnancy, Cancer of the Stomach and all conditions of the digestive organs where no food or even water can be retained. In Phthisis, it will increase weight

and strength far more rapidly than Cod Liver Oil.

KUMYSGEN is incomparable as a Food where easy digestion, high nutrition and palatability are desired.

A pound bottle of **KUMYSGEN** will be sent any Physician prepaid on receipt of fifty cents, which is about one-third its retail price. **KUMYSGEN** is now put up only in bottles holding 20 ounces and 5 pounds.

KUMYSGEN is much less expensive than ordinary Liquid Kumyss to prescribe and its keeping qualities are perfect, while the latter spoils in a very short time.

*KUMYSGEN, when first prepared, was not relished by some patients, but, as improved, commencing with batch 200, it will please the most delicate palate.

REED & CARNRICK, New York.

OUR PROPOSITION

DO YOU CARE TO CONSIDER IT?

Or do you prefer to grope along in the darkness, losing valuable time and wasting golden opportunities?

Never mind what the great Doctor Blank says of LONDONDERRY. You wish to establish your own opinions.

HOW? READ CAREFULLY!

We will furnish water free and pay all costs of chemical analysis of the urine for any clinical test, independently or in comparison with any water, natural or artificial, or any drug used as an anti-lithic!

We Claim that LONDONDERRY LITHIA will cure more cases of Rheumatism than any other water or any drug and stand ready for the test.

Londonderry Lithia Spring Water Co., Nashua, N. H.

NEW YORK:

CHICAGO: PHILADELPHIA:

76 Broad Street.

70 State Street. 1516 Chestnut Street.

MUIR & CO., 1516 Chestnut Street, Philadelphia,

Distributing Agents for Pennsylvania, Baltimore and Washington

ANTISEPTICS AND DISINFECTANTS.

The prevention of disease is the unselfish mission of the modern physician. Antiseptics and Disinfectants to-day occupy the first place in medical and surgical practice. We desire to call attention to the following Antiseptic and Disinfectant preparations:

Antiseptic Liquid arrests decomposition and destroys noxious gases that emanate from organic matter in sewers and elsewhere, and may be used in cellars,

barns, out-houses, and the sick-room.

Antiseptic Tablets are convenient for the extemporaneous preparation of antiseptic solutions of definite strength for disinfectant purposes and for antiseptic

sprays.

Eucalyptus and Thymol Antiseptic is adapted for use as an antiseptic internally, externally, hypodermatically, as a douche, a spray, by atomization, and as a deodorant. Its application in surgery is unlimited.

Sulphur Candles thoroughly employed are effectual in the fumigation and

disinfecting of rooms after infectious diseases.

Correspondence regarding any of our pure and accurate preparations of standard pharmaceuticals and specialties solicited.

PARKE, DAVIS & CO.

DETROIT, NEW YORK, KANSAS CITY, AND WALKERVILLE, ON

Mellin's Food For Infants and Invalids.

A SOLUBLE DRY EXTRACT of Barley Malt and Wheat, for addition to Fresh Cow's Milk.

Prepared upon the principles advanced by the eminent chemist, Baron Justus von Liebig.

EXPERIENCE—the supreme test—has absolutely proven that the best solution of the problem of infant feeding was made by Liebig, and that MELLIN'S FOOD prepared with milk is the nearest approximation to, and is the BEST SUBSTITUTE for, Mother's Milk which has ever been devised.

THE DOLIBER-GOODALE CO., BOSTON, MASS.

